

BIO-COX 60- salinomycin sodium powder
Huvepharma, Inc.

Bio-Cox 60

Front Panel

Bio-Cox®

Granular 60

Brand of Salinomycin Sodium

Type A Medicated Article

Net wt 50 LB (22.68 kg)

HUVEPHARMA®

Bag Back Panel - Mixing Directions and Permitted Claims

Bio-Cox®

Granular 60

Brand of Salinomycin Sodium

Type A Medicated Article

**For use in broiler, roaster, replacement
(breeder and layer) chickens, and quail only**

Active drug ingredient

Salinomycin equivalent to 60 grams salinomycin sodium activity per pound

For use in manufacturing medicated feeds only

Mixing directions

Thoroughly mix Bio-Cox 60 Granular with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium Activity Amount of Bio-Cox 60 Granular

(grams per ton)	Type A Medicated Article (60 g/lb) Per Ton of Feed	
	lb	grams
40	0.67	304
45	0.75	341
50	0.83	377
55	0.92	418
60	1.00	454

Feeding directions

Feed continuously as the sole ration

Important

Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

Restricted drug (California) - Use only as directed. Not for human use.

NADA #128-686, Approved by FDA

HUVEPHARMA®

☐ Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269

HUVEPHARMAA and Bio-Cox
are registered trademarks of
Huvepharma EOOD

Made in USA

Net wt 50 LB (22.68 kg)

P08-400310

Rev. 06-2016

Back Panel - Indications

Broiler, roaster and replacement (breeder and layer) chickens

Use Bio-Cox 60 Granular to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

Quail

Use Bio-Cox 60 Granular to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

Caution

May be fatal if accidentally fed to adult turkeys or to horses.

Warning


Do not feed to laying hens producing eggs for human consumption. When mixing and handling Bio-Cox 60 Granular premix, use protective clothing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Back Panel - Storage

Store below 25°C (77°F), excursions permitted to 37°C (99°F)

Bag Label Image

Bag Label Image



BIO-COX 60

salinomycin sodium powder

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:23243-9807
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALINOMYCIN SODIUM (UNII: 92UOD3BMEK) (SALINOMYCIN - UNII:62UXS86T64)	SALINOMYCIN SODIUM	60 g in 0.45 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-9807-5	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA128686	06/21/2017	

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (552691651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc - Van Buren		080346352	medicated animal feed manufacture, analysis, pack, label

Revised: 6/2017

Huvepharma, Inc.