

HUVEPHARMA



Net Contents: 500 mL (16.91 oz)
AGRIMYCIN® 200
(oxytetracycline injection)

NDC 23243-7430-6

NDC 23243-7430-6

AGRIMYCIN® 200 (oxytetracycline injection)

Antibiotic

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each mL contains 200 mg of oxytetracycline base as oxytetracycline dihydrate.

For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine

For animal use only

Approved by FDA under ANADA # 200-128

NET CONTENTS: 500 mL (16.91 oz)



HUVEPHARMA

NoCoating

AGRIMYCIN® 200 (oxytetracycline injection) is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline. Each mL contains 200 mg of oxytetracycline base as oxytetracycline dihydrate, and on a w/v basis, 40.0% 2-pyrrolidone, 5.0% povidone, 1.8% magnesium oxide, 0.2% sodium formaldehyde sulfoxylate (as a preservative), monoethanolamine and/or hydrochloric acid as required to adjust pH.

CAUTION: When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

Warnings: Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food. Rapid intravenous administration may result in animal collapse. Oxytetracycline should be administered intravenously slowly over a period of at least 5 minutes.

Precautions: Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL subcutaneously per injection site in adult beef and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

Refer to Package Insert for Complete Directions.

Not For Human Use

Storage: Store at room temperature 15°- 25°C (59°- 77°F). Keep from freezing. Use within 12 months of first puncture. Puncture a maximum of 100 times.

®Registered Trademark of Huvepharma, Inc.

TAKE TIME



OBSERVE LABEL DIRECTIONS

I-30035
0-3744-06
Rev. 08-2022

Manufactured for
Huvepharma, Inc.
Peachtree City, GA 30269

LOT NO.:

EXP. DATE:

NoCoating

No Coating

NDC 23243-7430-6

AGRIMYCIN® 200 (oxytetracycline injection)

Antibiotic

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each mL contains 200 mg of oxytetracycline base as oxytetracycline dihydrate.

For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine

For animal use only

Approved by FDA under ANADA # 200-128

NET CONTENTS: 500 mL (16.91 oz)



HUVEPHARMA

Cattle Dosage Guide

At the first signs of pneumonia or pinkeye,* administer a single dose of AGRIMYCIN® 200 (oxytetracycline injection) subcutaneously according to the following weight categories.**

Animal Weight (lb)	Number of mL or cc
100	4.5
200	9.0
300	13.5
400	18.0
500	22.5
600	27.0
700	31.5
800	36.0
900	40.5
1000	45.0
1100	49.5
1200	54.0

*See package insert for dosing instructions for other indicated diseases and full product information.

** Do not administer more than 10 mL at any one injection site (1-2 mL per site in small calves).

Discontinue treatment at least 28 days prior to slaughter.

Swine Dosage Guide

At the first signs of pneumonia,* administer AGRIMYCIN® 200 (oxytetracycline injection) by deep intramuscular injection according to the following weight categories.**

Animal Weight (lb)	Number of mL or cc
10	0.5
25	1.1
50	2.3
75	3.4
100	4.5
125	5.6
150	6.8
175	7.9
200	9.0
225	10.1
250	11.3
275	12.4
300	13.5
325	14.6

*See package insert for dosing instructions for other indicated diseases and full product information.

** Do not administer more than 5 mL at any one injection site.

Discontinue treatment at least 28 days prior to slaughter.

No Coating