



Product Data Sheet



Coxidin[®] 90 (monensin Type A medicated article)

Product Description

Coxidin[®] 90 (monensin Type A medicated article) is for the aid in the prevention of coccidiosis in poultry. Coxidin[®] 90 contains 90.7 grams of monensin per pound of premix.

Indications for Use

- **Broiler and Laying Hen Replacement Chicken and Layer Breeder Replacement Chicken:** as an aid in prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*.
- **Turkeys:** For the prevention of coccidiosis in turkeys caused by *Eimeria adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*.
- **Quail:** For the prevention of coccidiosis in growing Bobwhite quail caused by *Eimeria dispersa* and *E. lettyae*.

Withdrawal Information

No withdrawal period is required before slaughter when used according to label directions.

Storage and Packaging

Store at or below 25° C (77° F). Excursions permitted to 37° C (99° F). Coxidin[®] 90 is sold in 25 kg (55.12 lb) multiwall paper bags.

User Safety Warning

When mixing and handling Coxidin[®] 90, use protective clothing, impervious gloves, and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

IMPORTANT SAFETY INFORMATION: Coxidin[®] 90 (monensin Type A medicated article): Not for human use. Keep out of the reach of children. Do not feed undiluted. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. In the absence of coccidiosis in broiler chickens the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Not for broiler breeder replacement chickens. Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. To report adverse effects, access medical information, or obtain additional product information, call 1-877-994-4883. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>.



Coxidin® 90

(monensin
Type A medicated article)

For Use in Broiler Chicken, Laying Hen Replacement Chicken, Layer Breeder Replacement Chicken, Growing Turkey, and Growing Bobwhite Quail Feeds Only
Do not feed undiluted.

Net Weight **25 kg**
(55.12 lb)

Active Drug Ingredient – Monensin, USP - 90.7 g per pound.

Broiler and Laying Hen Replacement Chicken and Layer Breeder Replacement Chicken: As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

Turkeys: For the prevention of coccidiosis in turkeys caused by *Eimeria adenoeides*, *E. meleagrimitis* and *E. gallopavonis*.

Quail: For the prevention of coccidiosis in growing Bobwhite quail caused by *Eimeria dispersa* and *E. lettyae*.

Important: Must be thoroughly mixed in feeds before use. Mix thoroughly Coxidin 90 with a small quantity of non-medicated feed (e.g., 1 lb with 100-200 lb) before incorporation into the Type C medicated feed.

Mixing Directions (Broiler Chickens, Laying Hen Replacement Chickens, and Layer Breeder Replacement Chickens) – Thoroughly mix the following amounts of Coxidin 90 with a sufficient quantity of unmedicated feed to make one ton of complete chicken feed to provide 90 through 110 grams monensin per ton.

Monensin (Grams Per Ton)	Coxidin 90 (Lbs Per Ton of Type C Feed)
90	0.99
95	1.05
100	1.10
105	1.16
110	1.21

Feeding Directions – Feed continuously as the sole ration.

Mixing Directions (Turkeys) – Thoroughly mix the following amounts of Coxidin 90 with a sufficient quantity of unmedicated feed to make one ton of complete turkey feed to provide 54 through 90 grams monensin per ton.

Monensin (Grams Per Ton)	Coxidin 90 (Lbs Per Ton of Type C Feed)
54	0.60
66	0.73
78	0.86
90	0.99

Feeding Directions – For growing turkeys only, feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

Mixing Directions (Bobwhite Quail) – Thoroughly mix the following amounts of Coxidin 90 with a sufficient quantity of unmedicated feed to make one ton of complete quail feed to provide 73 grams monensin per ton.

Monensin (Grams Per Ton)	Coxidin 90 (Lbs Per Ton of Type C Feed)
73	0.80

Feeding Directions – For Bobwhite quail, feed continuously as the sole ration.

The Type C medicated feed must bear an expiry of 90 days from the date of manufacture.

Caution: Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. In the absence of coccidiosis in broiler chickens the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Not for broiler breeder replacement chickens. Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

NOT FOR HUMAN USE

Warning

Withdrawal Period: No withdrawal period is required when used according to labeling.

User Safety Warning: When mixing and handling Coxidin 90, use protective clothing, impervious gloves, and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Store at or below 25 °C (77 °F). Excursions permitted to 37 °C (99 °F).

Not to be used after the date printed on bag.

Restricted Drug (California) – Use Only as Directed

Approved by FDA under ANADA # 200-783

Huvepharma, Inc.

Peachtree City, GA 30269, USA

Questions/Comments? To report side effects, access medical information, or obtain additional product information, call Huvepharma, Inc. at 1-877-994-4883.

For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

COXIDIN is a registered trademark of Huvepharma EOOD.



KCP-S00180
Rev. 04-2024



Huvepharma, Inc. | 525 Westpark Drive, Suite 230 | Peachtree City, GA 30269
1-877-994-4883 | www.huvepharma.us

CoxidinPDS
Rev. 09-2024