


Client Information Sheet Dexafer-200

Dexafer-200 is a sterile Iron Dextran for Injection (200 mg/mL) supplied in 100-mL bottles for veterinary use only. **Dexafer-200** is not approved by the FDA.

FDA has agreed to allow Vetoquinol N.-A. Inc. to supply **Dexafer-200** in order to help address the anticipated shortage of Iron Dextran Injection, 200 mg/mL for prevention and treatment of iron deficiency anemia in newborn piglets in the U.S.A.

Dexafer-200 is approved by Health Canada and exported to the U.S.A. by Vetoquinol N.-A. Inc.

Dexafer-200 should be used according to the details of the Compendium text provided in the back of this Client Information Sheet. You can use **Dexafer-200** as you would use any other Iron Dextran for Injection in the prevention and treatment of iron deficiency in newborn piglets in the U.S.A. For example, site of administration as illustrated below.



ADVERSE DRUG EVENTS & PRODUCT DEFECTS

To report suspected adverse drug events and product defects, please contact Vetoquinol N.-A.'s toll-free number at 1-800-363-1700 and inform the representative that your call concerns the Canadian product, **Dexafer-200**. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae.

Vetoquinol N.-A. Inc.

2000, ch. Georges, Lavaltrie, QC, Canada J5T 3S5




Client Information Sheet Dexafer-200

Dexafer-200 is a sterile Iron Dextran for Injection (200 mg/mL) supplied in 100-mL bottles for veterinary use only. **Dexafer-200** is not approved by the FDA.

FDA has agreed to allow Vetoquinol N.-A. Inc. to supply **Dexafer-200** in order to help address the anticipated shortage of Iron Dextran Injection, 200 mg/mL for prevention and treatment of iron deficiency anemia in newborn piglets in the U.S.A.

Dexafer-200 is approved by Health Canada and exported to the U.S.A. by Vetoquinol N.-A. Inc.

Dexafer-200 should be used according to the details of the Compendium text provided in the back of this Client Information Sheet. You can use **Dexafer-200** as you would use any other Iron Dextran for Injection in the prevention and treatment of iron deficiency in newborn piglets in the U.S.A. For example, site of administration as illustrated below.



ADVERSE DRUG EVENTS & PRODUCT DEFECTS

To report suspected adverse drug events and product defects, please contact Vetoquinol N.-A.'s toll-free number at 1-800-363-1700 and inform the representative that your call concerns the Canadian product, **Dexafer-200**. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae.

Vetoquinol N.-A. Inc.

2000, ch. Georges, Lavaltrie, QC, Canada J5T 3S5



Dexafer-200

Vetoquinol

STERILE IRON DEXTRAN 20% W/V SOLUTION FOR INJECTION, Mfr. Std.

Veterinary Use Only

DIN 02252724

ACTIVE INGREDIENT: Each mL contains the equivalent of 200 mg of iron as ferric hydroxide dextran complex.

Preservative: Phenol 0.5% W/V

INDICATIONS: For the prevention and treatment of iron deficiency anaemia in new-born piglets.

DIRECTIONS FOR USE: For the prevention and treatment of iron deficiency anaemia in baby pigs, inject 1 mL (200 mg iron) intramuscularly, preferably in the neck region. Disinfect the site of injection with iodine or alcohol before injecting.

CAUTIONS: If symptoms of iron deficiency persist, consult a veterinarian.

STORAGE: Store at room temperature (between 15°C and 25°C). Avoid freezing.

Vetoquinol N.-A. Inc.

2000, ch. Georges, Lavaltrie, QC, Canada J5T 3S5



Dexafer-200

Vetoquinol

STERILE IRON DEXTRAN 20% W/V SOLUTION FOR INJECTION, Mfr. Std.

Veterinary Use Only

DIN 02252724

ACTIVE INGREDIENT: Each mL contains the equivalent of 200 mg of iron as ferric hydroxide dextran complex.

Preservative: Phenol 0.5% W/V

INDICATIONS: For the prevention and treatment of iron deficiency anaemia in new-born piglets.

DIRECTIONS FOR USE: For the prevention and treatment of iron deficiency anaemia in baby pigs, inject 1 mL (200 mg iron) intramuscularly, preferably in the neck region. Disinfect the site of injection with iodine or alcohol before injecting.

CAUTIONS: If symptoms of iron deficiency persist, consult a veterinarian.

STORAGE: Store at room temperature (between 15°C and 25°C). Avoid freezing.

Vetoquinol N.-A. Inc.

2000, ch. Georges, Lavaltrie, QC, Canada J5T 3S5

