

**Residue Warnings:**

Exceeding the daily dosage of 3,000 units per pound of body weight, administering for more than four consecutive days, or exceeding the maximum injection site volume per injection site may result in antibiotic residues beyond the withdrawal time. Milk taken from treated dairy animals within 48 hours after the last treatment must not be used for food. Discontinue use of this drug for the following time period before treated animals are slaughtered for food:

Cattle - 14 days, Sheep - 9 days, Swine - 7 days.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Warning:**

Do not use in horses intended for human consumption. Not for use in humans. Keep out of reach of children.

3



Lot:

Exp:

0

003107L0X

# AGRI-CILLIN®

(penicillin G procaine injectable suspension)

NDC 57561 065 02

300,000 UNITS PER mL

**ANTIBIOTIC**  
FOR INTRAMUSCULAR INJECTION ONLY  
NADA 065-010, APPROVED BY FDA

Net Contents: 100 mL

**AgriLabs®**

Making A Healthy Difference®

Read package insert for complete product information before using the drug.

**INDICATIONS**

Agri-Cillin is indicated for treatment of bacterial pneumonia (shipping fever) caused by *Pasteurella multocida* in cattle and sheep, erysipelas caused by *Erysipelothrix rhusiopathiae* in swine, and strangles caused by *Streptococcus equi* in horses.

**DESCRIPTION**

Each mL contains 300,000 units Penicillin G Potassium\*, 139.0 mg Procaine Hydrochloride\*, 2.0% Procaine Hydrochloride, 3.0 mg Potassium Phosphate Monobasic, 6.0 mg Potassium Phosphate Dibasic, 0.4% Sodium Formaldehyde Sulfoxylate, 0.4 mg Polysorbate 80, 2.0 mg Lecithin, 0.1% Methyl Paraben, 0.01% Propyl Paraben, 0.15% Sodium Carboxymethylcellulose, q.s. Water for Injection.

\* Penicillin G Potassium and Procaine Hydrochloride react to form penicillin G procaine.

**DOSAGE**

Agri-Cillin should be administered by the intramuscular route. The product is ready for injection after warming the vial to room temperature and shaking to ensure a uniform suspension. The recommended daily dosage of penicillin is 3,000 units per pound of bodyweight (one mL per 100 lbs bodyweight). Continue daily treatment until recovery is apparent and for at least one day after symptoms disappear, usually in two to three days. Treatment should not exceed four consecutive days. No more than 10 mL should be injected at any one site in adult livestock: rotate injection sites for each succeeding treatment.

**Restricted Drug - California. Use Only as Directed.**

Store at 2-8°C (36-46°F).  
SHAKE WELL BEFORE USING.

Made in the UK.  
Norbrook Laboratories Limited  
Newry, Northern Ireland

**TAKE TIME**

Rev 0217

**OBSERVE LABEL DIRECTIONS**