



# AGRI-MECTIN<sup>®</sup> (ivermectin) POUR-ON FOR CATTLE

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

#### INDICATIONS:

For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

See package insert for complete indications and use directions.

#### DOSAGE:

The dose rate is 1 mL for each 22 lb of bodyweight. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

#### ADMINISTRATION:

Because of the solvents used in Agri-Mectin<sup>®</sup> Pour-On, the applicator gun from Simcro Tech, or equivalent, is recommended. Other applicators may exhibit compatibility problems resulting in locking, incorrect dosage or leakage. Remove the shipping cap from the backpack container and replace with the vent cap provided. Attach the hose from the automatic dosing equipment to the outlet from the vent cap. Follow the applicator gun manufacturer's directions for priming the gun, adjusting the dose, and care of the applicator gun following use.

**Restricted Drug - California. Use Only as Directed.**

© Registered Trademark of Agri Laboratories, Ltd.

Manufactured for:  
Agri Laboratories, Ltd.,  
St. Joseph, MO 64503



3 57561 00223 4



**AgriLabs<sup>®</sup>**  
Making A Healthy Difference<sup>®</sup>

6214102C04

## AGRI-MECTIN<sup>®</sup> (ivermectin) POUR-ON FOR CATTLE

This topically applied formulation of Agri-Mectin<sup>®</sup> (ivermectin) delivers internal and external parasite control in one convenient low-volume application.

#### PRECAUTIONS:

**Store away from excessive heat (104°F/40°C) and protect from light.**

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced.

Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when Agri-Mectin<sup>®</sup> (ivermectin) Pour-On is stored at temperatures below 32°F. Allowing it to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g. caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Agri-Mectin<sup>®</sup> Pour-On is not recommended for use in species other than cattle.

Lot No:  
Exp Date:

# AGRI-MECTIN<sup>®</sup> (ivermectin)

## POUR-ON FOR CATTLE

## AGRI-MECTIN<sup>®</sup> (ivermectin) POUR-ON FOR CATTLE

**Contains: 5 mg ivermectin/mL**

ANADA 200-272, Approved by FDA

### PARASITICIDE

**Kills:** Roundworms, (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies

**Contains: 200 x 550 lb Doses**

**Net Contents: 169 fl oz (5 L)**



**AgriLabs<sup>®</sup>**  
Making A Healthy Difference<sup>®</sup>

## AGRI-MECTIN<sup>®</sup> (ivermectin) POUR-ON FOR CATTLE



#### ANIMAL SAFETY:

Studies conducted in the U.S.A. have demonstrated the safety margin for ivermectin. Based on plasma levels, the topically applied formulation is expected to be at least as well tolerated by breeding animals as is the subcutaneous formulation which had no effect on breeding performance.

**RESIDUE INFORMATION:** Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

#### WARNING! NOT FOR USE IN HUMANS

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

#### Disposal:

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

**Keep this and all drugs out of the reach of children.**

**WARNING! FLAMMABLE!  
KEEP AWAY FROM HEAT,  
SPARKS, OPEN FLAME,  
AND OTHER SOURCES OF IGNITION.**