

ALBADRY PLUS[®]

Suspension

PENICILLIN G PROCAINE AND NOVIOBIOCIN SODIUM

DRY COW FORMULA



Indicated for intramammary infusion in the treatment of subclinical mastitis caused by susceptible strains of *Staphylococcus aureus* and *Streptococcus agalactiae*.

Contains: 400 mg novobiocin sodium plus 200,000 IU penicillin G procaine per 10 mL syringe

1404000
006544L03

NADA #55-098, Approved by FDA
144-10 mL (.33 Fl Oz) PLASTET[®] Disposable Syringes / 144 ALCOHOL PADS INCLUDED



2504

zoetis 

READ RESIDUE WARNING

DRY



DISCARD MILK FOR
72 hrs
AFTER CALVING



DO NOT SLAUGHTER COW FOR
30 days
AFTER LAST TREATMENT

RESTRICTED DRUG—USE ONLY AS DIRECTED (California)
FOR USE IN ANIMALS ONLY—NOT FOR HUMAN USE

STORAGE CONDITIONS

Store at controlled room temperature 20° to 25°C (68° to 77° F).

PRECAUTION: Administration of this product in any manner other than shown under **DOSAGE** may result in drug residues.

WARNINGS

- 1) Do not use less than 30 days prior to calving.
- 2) Milk from treated cows must not be used for food during the first 72 hours after calving.
- 3) Treated animals must not be slaughtered for food for 30 days following udder infusion.

ADMINISTRATION

At the time of drying off, but not less than 30 days prior to calving, milk the udder dry. Wash the teats and udder thoroughly with warm water containing a suitable dairy antiseptic. Dry the teats and udder thoroughly. Infuse each quarter using the following procedure. Using the alcohol pads provided, scrub each teat and clean using a separate pad for each teat. Warm ALBADRY PLUS Suspension (penicillin G procaine and novobiocin sodium) to body temperature and shake thoroughly. Remove cap from tip of PLASTET and insert tip into the teat canal. Instill entire contents of PLASTET into the quarter. Massage the udder after treatment to distribute the ALBADRY PLUS Suspension throughout the quarters. Using a suitable teat dip, dip all teats following treatment.

INDICATIONS FOR USE

ALBADRY PLUS is indicated for the treatment, in dry cows only, of subclinical mastitis caused by susceptible strains of *Staphylococcus aureus* and *Streptococcus agalactiae*.

DOSAGE

Infuse one tube per quarter at start of drying period (but not less than 30 days prior to calving). See package insert for complete product information.

DESCRIPTION

Each 10 mL syringe contains:

Novobiocin sodium equiv. to novobiocin.....	400 mg
Penicillin G procaine.....	200,000 IU
Chlorobutanol anhydrous	
(Ethanol derivative—used as a preservative).....	50 mg


in a special blend vehicle.

Manufactured by a non-sterilizing process.

SHAKE WELL BEFORE USING

Made in the United Kingdom for
Zoetis Inc. • Kalamazoo, Michigan 49001 USA
By Norbrook Laboratories Limited • Newry, BT33 6JR Northern Ireland

zoetis
www.zoetis.com



0009313907 7