

### FOR ANIMAL USE ONLY

To prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use ½ pint (8 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of AmproMed™ P FOR POULTRY 9.6% Oral Solution.

### PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:

For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

TO PREPARE DOSAGE LEVELS OF	AmproMed™ P FOR POULTRY 9.6% Oral Solution PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

**Note:** Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

**Benzoic acid as Sodium Benzoate 0.1% added as preservative.**

**STORAGE:** Store container in an upright position. Store between 5°C-25°C (41°F-77°F) with brief excursions to 40°C.

To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

AmproMed is a Trademark of Bimeda, Inc.



## AmproMed™ P FOR POULTRY (amprolium)

9.6% Oral Solution

Coccidiostat



Water-soluble treatment for coccidiosis  
Treats 800 Gallons at 0.012% level

Restricted Drug (California) – Use Only as Directed

ANADA 200-496, Approved by FDA



Net Contents:  
1 Gallon (128 fl oz) (3.785 L)

ACTIVE INGREDIENT: amprolium..9.6%

Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058  
www.bimeda.com

**INDICATIONS:** AmproMed™ P FOR POULTRY 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**USE DIRECTIONS:** Give amprolium at the 0.012% level (8 fl oz AmproMed™ P FOR POULTRY 9.6% Oral Solution per 50 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

**WARNING:** Keep this and all drugs out of reach of children. NOT FOR HUMAN USE.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

Product No. 1AMP024 8AMP016 Rev. 06/12



Lot:  
Exp:

8.0"

4.0"

# AmproMed™ P FOR POULTRY

*(amprolium)*

**9.6% Oral Solution/Coccidiostat**

ANADA 200-496, Approved by FDA

## INDICATIONS

AmproMed™ P FOR POULTRY 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens.

## BENEFITS

- **Effective:** breaks up the coccidia life cycle
- **Economical:** one gallon of concentrate treats 800 gallons of drinking water at a 0.012% level
- **Convenient:** works in automatic drinking water proportioners to treat the entire flock
- **Easy to mix:** concentrated solution disperses into the drinking water
- **Palatable:** medicated drinking water readily accepted by birds
- **Safe:** approved by FDA



## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1AMP024	1 gallon (128 fl oz) (3.785 L)	4 x 1 gallon (512 fl oz) (15.141 L)

See reverse side for Administration and Dosage.

### Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: [sales@bimedaus.com](mailto:sales@bimedaus.com)



# AmproMed™ P FOR POULTRY

(amprolium)

9.6% Oral Solution/Coccidiostat

ANADA 200-496, Approved by FDA

**FOR ORAL USE IN ANIMALS ONLY**  
**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

Restricted Drug (California) – Use Only as Directed

Water-soluble treatment for coccidiosis  
 Treats 800 Gallons at 0.012% level

**ACTIVE INGREDIENT:**

Amprolium.....9.6%

**INDICATIONS:**

AmproMed™ P FOR POULTRY 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

**USE DIRECTIONS:**

Give amprolium at the 0.012% level (8 fl oz AmproMed™ P FOR POULTRY 9.6% Oral Solution per 50 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

**To prepare 50 Gallons of Medicated Water**

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use ½ pint (8 fluid ounces) of AmproMed™ P FOR POULTRY 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of AmproMed™ P FOR POULTRY 9.6% Oral Solution.

**PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:**

For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

TO PREPARE DOSAGE LEVELS OF	AmproMed™ P FOR POULTRY 9.6% Oral Solution PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

**Note:** Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

**Benzoic acid as Sodium Benzoate 0.1% added as preservative.**

**WARNING:**

Keep this and all drugs out of reach of children. NOT FOR HUMAN USE.

**PRECAUTIONS:**

FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

**STORAGE:**

Store container in an upright position. Store between 5°C-25°C (41°F-77°F) with brief excursions to 40° C.

To obtain an MSDS or for assistance contact Bimeda, Inc. at 1-888-524-6332.

Manufactured by:  
 Bimeda, Inc.  
 Le Sueur, MN 56058  
 8AMP016

