

NADA 12-350
Approved by the FDA

Product 956150

AMPROL® 25%

(amprolium)

Type A Medicated Article

AMPROL (amprolium) 25% is intended for use in (1) preventing outbreaks of coccidiosis in growing chickens, turkeys and pheasants and (2) development of active immunity to coccidiosis in replacement chickens. AMPROL 25% is also intended for use in the prevention and treatment of coccidiosis in laying chickens.

ACTIVE DRUG INGREDIENT: Amprolium..... 25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

STORAGE: Store at or below 25°C (77°F), excursions permitted to 40°C (104°F). Avoid prolonged exposure to high humidity.

SEE DIRECTIONS ON BACK PANEL

Manufactured by:
HUVEPHARMA, INC. St. Louis, MO 63116
Distributed by:
HUVEPHARMA, INC. Peachtree City, GA 30269
For questions, call 1-877-4AMPROL

Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269



® AMPROL is a registered trademark of Huvepharma, Inc.

Net Wt.: 50 lb (22.68 kg)



WARNING

Keep this and all drugs out of the reach of children. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, obtain an MSDS or for assistance, contact Huvepharma at 1-877-4AMPROL.

MIXING DIRECTIONS

AMPROL® (amprolium) 25% should be thoroughly and evenly mixed in the feed. AMPROL 25% may be used to manufacture a Type C medicated feed in the concentration range of 0.004% to 0.025%. The table below shows the amount of AMPROL 25% to be used in each ton (2000 lb) of feed to obtain desired levels of amprolium in the Type C medicated feed.

To aid in even distribution of AMPROL 25% in the finished feed, prepare a mixture of AMPROL 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and mix thoroughly.

Feeding Level of Amprolium in Type C Medicated Feed	Use This Amount of AMPROL 25% Per Ton
0.004%	5 oz
0.006%	8 oz
0.008%	10 oz
0.0125%	1 lb
0.0175%	1 lb 6 1/2 oz
0.025%	2 lb

AMOUNT AND DIRECTIONS FOR USE

CHICKENS

Broilers: Prevention: Use 0.0125% amprolium Type C medicated feed continuously for most field conditions as they exist under modern management practices. Where severe coccidiosis conditions exist and where immunity is not required, use 0.025% amprolium Type C medicated feed. For field conditions where only *E. tenella* is the major problem, use 0.008%-0.0125% amprolium Type C medicated feed. Increasing levels of amprolium may not prevent coccidiosis caused by strains of *Eimeria* commonly found to be resistant to Amprolium.

Replacements: Immunity Development or Prevention
Prevention Program: Where immunity development is not desirable, use 0.0125%-0.025% amprolium Type C medicated feed continuously from day old until onset of production.

Immunity Development Program: Use 0.004%-0.0125% amprolium Type C medicated feed continuously until onset of production.

Selection of the level to be used should be based upon comparative hazard of infection with cecal and intestinal species. The higher levels will interfere with the development of immunity to *E. tenella* (cecal). The following are suggested feeding schedules for various conditions of coccidial exposure. The planning and evaluation of any program should be in the hands of a veterinarian or poultry pathologist who is familiar with the specific operation and with the general nature of disease problems in the area.

1. Severe exposure	2. Moderate exposure	3. Slight exposure
0-5 weeks of age 0.0125%	0-5 weeks of age 0.008%-0.0125%	0-5 weeks of age 0.004%-0.0125%
5-8 weeks of age 0.008%-0.0125%	5-8 weeks of age 0.006%-0.0125%	5-8 weeks of age 0.004%-0.0125%
over 8 weeks of age 0.004%-0.0125%	over 8 weeks of age 0.004%-0.0125%	over 8 weeks of age 0.004%-0.0125%

LAYING CHICKENS: Treatment: Use 0.025% amprolium Type C medicated feed for two weeks in severe outbreaks or 0.0125% amprolium Type C medicated feed for two weeks in moderate outbreaks.
Prevention: Use 0.0125% amprolium Type C medicated feed continuously.

TURKEYS: Prevention: Use 0.0125%-0.025% amprolium Type C medicated feed continuously.

PHEASANTS: For the prevention of coccidiosis in growing pheasants caused by *Eimeria colchici*, *E. duodenalis*, and *E. phasiani*. Type C medicated feed with 0.0175% amprolium should be fed continuously as the sole ration.

CAUTION

Use as the sole source of amprolium. Do not change the litter while giving this feed unless absolutely necessary. If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.

Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

In replacement flocks the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.

Fertility, hatchability and other reproductive data are not available on amprolium in breeding pheasants.

Do not use in feeds containing bentonite. Restricted Drug (California) - Use Only as Directed. Not for Human Use.

AMPROL® 25%
(amprolium)
Type A Medicated Article



AMPROL® 25%
(amprolium)
Type A Medicated Article



LOT NO & EXP DATE

120733 6/13
FREIGHT SHIPPING BAG
MEETING REQUIREMENTS OF
APPLICABLE FREIGHT CLASSIFICATION
QUANTITIES BY
BANCROFT BAG, INC.
WEST MADISON, LOUISIANA
PO1625 B-1

AMPROL® 25%

(amprolium)

Type A Medicated Article

5046-03
Rev 11-2013