

Avatec[®]

Lasalocid Type A Medicated Article Coccidiocidal

Active drug ingredient

Each pound contains 90.7 grams (20%) of lasalocid (as lasalocid sodium activity) in a carrier suitable for incorporation in feed.

For use in the manufacture of medicated feeds only

For the prevention of coccidiosis in

Target Species:

Caused by:

Broiler or Fryer Chickens:

Eimeria tenella, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*

Growing Turkeys:

E. meleagrimitis, *E. gallopavonis*, *E. adenoides*

Rabbits:

Eimeria stiedae

Chukar Partridges:

Eimeria legionensis

Directions for use

Broiler, Fryer Chickens and Growing Turkeys

Use Avatec[®] continuously in the feed to provide from 68 g per ton (0.0075%) to 113 g per ton (0.0125%) lasalocid concentration. The dosage range allows adjustment of drug level to severity of exposure. The higher levels are indicated for severe exposure. Consult a poultry disease diagnostician for advice regarding the optimal level of drug.

Rabbits

Use Avatec[®] continuously in the feed for rabbits up to 6 1/2 weeks of age to provide 113 g per ton (0.0125%) lasalocid concentration.

Chukar Partridges

Use Avatec[®] continuously in the feed for young birds up to 8 weeks of age to provide 113 g per ton (0.0125%) lasalocid concentration.

Mixing directions

Thoroughly mix the following amounts of Avatec[®] to make one ton of feed to provide from 68 g per ton (0.0075%) to 113 g per ton (0.0125%) lasalocid concentration. When adding less than one pound of Type A medicated article per ton of feed, it is recommended that the Type A medicated article first be mixed with a small amount of feed (10-25 lbs.) prior to incorporation into the final feed.

Dosage for use in target species	Concentration % in Feed	Lasalocid grams/ton	lbs Avatec [®] per ton feed
Broiler, Fryer Chickens and Growing Turkeys	0.0075%	68	3/4
	0.0100%	90.7	1
	0.01125%	102	1 1/8
Broiler, Fryer Chickens, Growing Turkeys, Rabbits, Chukar Partridges	0.0125%	113	1 1/4

Warning

When mixing and handling lasalocid Type A medicated article, use protective clothing, impervious gloves and a dust mask. Avoid contact with eyes. Operators should wash thoroughly with soap and water after handling.

Note: In addition to label statements of identity, ingredients and directions for use, labeling of feeds containing Avatec[®] must contain the following statements:

FOR BROILER OR FRYER CHICKENS ONLY and/or
FOR GROWING TURKEYS ONLY and/or
FOR CHUKAR PARTRIDGES (UP TO EIGHT WEEKS OF AGE) ONLY and/or
FOR RABBITS (UP TO SIX AND ONE HALF WEEKS OF AGE) ONLY

Restricted drug (California) - Use only as directed.

Not for human use.

Additionally, **all** feed labeling **must** state:
FEED CONTINUOUSLY AS THE SOLE RATION

Store below 25°C (77°F), excursions permitted to 37°C (99°F).

NADA 96-298, Approved by FDA

zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007



Net wt 50 LB (22.68 kg)

Take Time



Observe Label Directions

Made in USA

10005700