

RESIDUE WARNINGS: Cattle must not be slaughtered for human consumption within 4 days of the last treatment. Milk that has been taken during treatment and for 36 hours after the last treatment must not be used for food. Not for use in dry dairy cows. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal. Not for use in horses intended for food. Approved only for intravenous administration in cattle. Intramuscular administration has resulted in violative residues in the edible tissues of cattle sent to slaughter.

Store at or below 25°C (77°F) Do not Freeze
Before using this drug, read package insert for complete product information.

NDC 0061-0851-03
Sterile

100 mL
Multiple-Dose Vial
50 mg/mL

Banamine®

(flunixin meglumine injection)

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
NADA #101-479, Approved by FDA.




COPY

Only for Intravenous Use in Beef and Dairy Cattle. Not for Use in Dry Dairy Cows and Veal Calves. For Intravenous and Intramuscular Use in Horses.

USE WITHIN 28 DAYS OF FIRST PUNCTURE AND PUNCTURE A MAXIMUM OF 10 TIMES, WHEN USING A DRAW-OFF SPIKE OR NEEDLE WITH BORE DIAMETER LARGER THAN 18 GAUGE. DISCARD ANY PRODUCT REMAINING IN THE VIAL IMMEDIATELY AFTER USE.

Distributed by:
Intervet Inc. d/b/a/Merck Animal Health
Madison, NJ 07940
Copyright © 2011
Intervet Inc., a subsidiary of Merck and Co., Inc.
All rights reserved. Made in Germany Rev. 01/17

197769 R10



LOT
EXP

Varnish and
Ink free