Artwork Date: March 21, 2016 V9
Item Size: 5.50" x 14.92" screen roll stock
Scale: 100%
File Name: 8BIL014 Bilovet 100 g POU.ai

Design/Packaging/Production

CMYK	
CMYK + SPOT	
SPOT COLOURS (Pantone)	
PMS	Black
PMS	7511
PMS	1805
PMS	

Labels						Cartons					
No Varnish Area	Left or Right	Unwind #	1	2	3	4	Dieline Verified	Yes or No	Correct Lot/Exp position	Yes or No	New Carton: Mock-Up Approved	Yes or No
Numbered on Back	Yes or No	Sign/Date	Sign/Date	Sign/Date	Sign/Date							

Department	Signature	Date	NO	YES
Sales & Marketing				
Packaging / Production				
QC				
Regulatory - MTC				
Regulatory - U.S.				
Customer				



Bilovet™ (tylosin tartrate) Soluble Powder

FOR ORAL USE IN CHICKENS, TURKEYS, SWINE, AND HONEY BEES

DOSAGE AND ADMINISTRATION

Dosages:
Chickens: CRD indications: 2,000 mg/gallon (528 ppm) in drinking water.
Turkeys: 2,000 mg/gallon (528 ppm) in drinking water.
Swine: 250 mg/gallon (66 ppm) in drinking water.
Honey Bees: 200 mg/colony in confectioners/powdered sugar.

Mixing Directions for Medicated Drinking Water:
Always add the water to the powder.
Do not pour the powder into the water. Prepare a fresh Bilovet™ Soluble Powder solution every three days. When mixing and handling tylosin, use protective clothing and impervious gloves. If using a water medicating pump see table below, otherwise mix as follows:
To assure thorough dissolution, first place the contents of one packet in a mixing container and add one gallon of water (3785 mL) to the powder to make a concentrated solution. To make medicated drinking water containing 250 mg/gallon (66 ppm), mix this concentrated solution with water to make 400 gallons (1514 liters) of medicated drinking water.
To make medicated drinking water containing 2,000 mg/gallon (528 ppm), mix this concentrated solution with water to make 50 gallons (189 liters) of medicated drinking water.

Mixing Directions for Water Medicating Pump (1:128 inclusion)*:

Desired Concentration in Drinking Water	Packets of Bilovet™	Volume of Water to Make Stock Solution
250 mg/gallon (66 ppm)	1	3 gallons + 13 ounces
2,000 mg/gallon (528 ppm)	10	3 gallons + 115 ounces

*This table applies only if the water medicating pump is set to deliver 1 ounce of stock solution per gallon of drinking water.

Mixing Directions for use in Honey Bees: Mix 200 mg tylosin in 20 g confectioners/powdered sugar. Use immediately.

DIRECTIONS FOR USE
Chickens: CRD indications: Administer medicated drinking water for three days; however, medicated water may be administered for one to five days depending upon severity of infection. Treated chickens must consume enough medicated water to provide 50 mg per pound of body weight per day. Only medicated water should be available to the birds.
Turkeys: Administer medicated drinking water for three days; however, medicated water may be administered for two to five days depending upon severity of infection. Treated turkeys must consume enough medicated water to provide 60 mg per pound of body weight per day. Only medicated water should be available to the birds.
Swine: SD indication: Administer medicated drinking water for 3 to 10 days, depending upon severity of infection. Alternatively, administer medicated drinking water for 3 to 10 days, followed by 40 to 100 g of tylosin per ton of complete feed (Type C medicated feed manufactured from tylosin phosphate Type A medicated article) for 2 to 6 weeks. Only medicated water should be available to swine while medicating with Bilovet™ Soluble Powder. **PPE indication:** Administer medicated drinking water for 3 to 10 days, followed by 40 to 100 g of tylosin per ton of complete feed (Type C medicated feed manufactured from tylosin phosphate Type A medicated article) for 2 to 6 weeks. Only medicated water should be available to swine while medicating with Bilovet™ Soluble Powder.
Honey Bee Colonies: Administer three treatments of medicated confectioners sugar once weekly for 3 weeks. The 200 mg dose is applied (dusted) over the top bars of the brood chamber.

WARNINGS
User Safety Warnings: Not for Use in Humans. Keep out of Reach of Children. Avoid contact with human skin. Exposure to tylosin may cause a rash.
Residue Warnings: Chickens must not be slaughtered for food within 24 hours after treatment. Do not use in layers producing eggs for human consumption. **Turkeys** must not be slaughtered for food within five days after treatment. **Swine** must not be slaughtered for food within 48 hours after treatment. **Honey Bees:** The drug should be fed early in the spring or fall and consumed by the bees before the main honey flow begins, to avoid contamination of production honey. Complete treatments at least 4 weeks prior to main honey flow.
Store at or below 25°C (77°F). Excursions Permitted to 40°C (104°F). Avoid Moisture.
To report suspected adverse events, for technical assistance, or to obtain a Material Safety Data Sheet (MSDS), call 1-888-524-6332.

UPC Code
8-998855-30090-4

TAKE TIME
OBSERVE LABEL DIRECTIONS

Bimeda

Bilovet™ is a Trademark of Bimeda, Inc.
N.A. Corp. Address: Bimeda Inc., One Tower Lane, Oakbrook Terrace, IL 60181

Product No: 1BIL004
8BIL014 Rev. 03/16