

NO COPY

INDICATIONS AND DIRECTIONS FOR USE (cont'd)
Chlortetracycline hydrochloride - 102.4 g per packet (64 g/lb)

GROWING TURKEYS	DOSE
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline	400 mg/gallon (2 packets per 512 gallons). This is equivalent to 1/2 oz of powder per 5 gallons of water. Mix until dissolved. Administer at this rate in the total water consumed over a full, 24-hour period. Administer 7 to 14 days.
Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis or coronaviral enteritis) susceptible to chlortetracycline	25 mg/lb body weight/day (1 packet for every 4,096 lbs of turkeys.) 1/8 oz treats a 20 lb turkey for one day. Mix until dissolved. Administer in the total water consumed over a full, 24-hour period. Administer to 14 days.
SWINE	DOSE - 10 mg/lb body weight/day
Control and treatment of bacterial enteritis (scours) caused by <i>Escherichia coli</i> and <i>Salmonella</i> spp. and bacterial pneumonia associated with <i>Pasteurella</i> spp., <i>Actinobacillus pleuropneumoniae</i> spp. and <i>Klebsiella</i> spp. susceptible to chlortetracycline	(5 packets per 512 gallons will treat 512 100-lb pigs per day) 1/4 oz of powder will treat a 100 lb pig for one day. Mix until dissolved. Administer at this rate in the total water consumed over a full, 24-hour period. Administer for 3 to 5 days.
CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE	DOSE - 10 mg/lb body weight/day
Control and treatment of bacterial enteritis (scours) caused by <i>Escherichia coli</i> and <i>Salmonella</i> spp. and bacterial pneumonia associated with <i>Pasteurella</i> spp., <i>Histophilus</i> spp. and <i>Klebsiella</i> spp. susceptible to chlortetracycline	Use 1/2 oz of powder for each 200 lbs of calf weight, per day. Mix until dissolved. Administer at this rate in the total water consumed over a full, 24-hour period, or as a drench in divided doses. Administer for 3 to 5 days.

Water Medicators: Dissolve the contents of one 25.6 oz packet in 100 oz of fresh water to prepare a stock solution, that when metered at the rate of 1 oz per gallon of drinking water, will deliver 1000 mg of chlortetracycline HCl per gallon. At this rate, a gallon of drinking water will treat a 100 lb pig or a 100 lb calf at the rate of 10 mg per lb of body weight for one day. Adjust if needed based on actual water intake to achieve 10 mg per lb of body weight.

CORNER RADIUS

Chlortetracycline Soluble Powder Concentrate

(chlortetracycline hydrochloride)
Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

NO COPY

NO COPY

For Animal Use in Drinking Water

Chlortetracycline hydrochloride 64 g per pound

This 25.6 oz packet contains 102.4 g chlortetracycline HCl and will make the resulting concentrations:

- 1024 gallons containing 100 mg chlortetracycline HCl per gallon
- 512 gallons containing 200 mg chlortetracycline HCl per gallon
- 256 gallons containing 400 mg chlortetracycline HCl per gallon
- 102.4 gallons containing 1000 mg chlortetracycline HCl per gallon

Each teaspoon of Chlortetracycline Soluble Powder Concentrate contains 500 mg of chlortetracycline HCl

Store at controlled room temperature, 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).

Not for human use.

HUVEPHARMA is a registered trademark of Huvepharma EOOD.



Net Wt. 25.6 oz (725.7 g)

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269

KCP-N00007
Rev. 05-2017

Restricted Drug (California) - Use only as directed.
ANADA 200-441, Approved by FDA

CORNER RADIUS

GENERAL INFORMATION ON DOSAGE

Dosages in terms of packets per 512 gallons are based on stated dosages per unit of body weight and average water consumption of the species. Weather conditions, ambient temperature, humidity, age, class of livestock and other factors may affect consumption and, except where calves are drenched, the unit dosage should be used as a guide to effective use in drinking water. Animal must actually consume enough medicated water to provide the desired therapeutic dose under the conditions that prevail.

PRECAUTIONS

Use as the sole source of chlortetracycline. Prepare fresh solutions daily. When used in plastic or stainless steel waterers or automatic waterers,

prepare fresh solutions every 24 hours. This product has not been demonstrated to provide adequate stability in galvanized waterers. Stability loss may reduce effectiveness of the product. Do not administer in milk or milk replacers. When feeding milk or milk replacers, administration one hour before or two hours after feeding is recommended.

RESIDUE WARNINGS

Do not administer to chickens, growing turkeys, swine and cattle within 24 hours of slaughter. Do not use in lactating dairy cattle and laying chickens. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

INDICATIONS AND DIRECTIONS FOR USE

Medicate chickens and turkeys continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. The dosage ranges permitted provide for different levels based on the severity of the infection. If improvement is not noted in 24 - 48 hours, consult your veterinarian.

CHICKENS	DOSE
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline.	200 - 400 mg/gallon (1 - 2 packets per 512 gallons). This is equivalent to 1/4 - 1/2 oz of powder per 5 gallons of water. Mix until dissolved. Administer at the indicated rates in the total water consumed over a full, 24-hour period. Administer 7 to 14 days.
Control of Chronic Respiratory Disease (CRD) and air-sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline.	400 - 800 mg/gallon (2 - 4 packets per 512 gallons). This is equivalent to 1/2 - 1 oz of powder per 5 gallons of water. Mix until dissolved. Administer at the indicated rates in the total water consumed over a full, 24-hour period. Administer 7 to 14 days.
Control of mortality due to fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline.	1000 mg/gallon (5 packets per 512 gallons). This is equivalent to 1/4 oz of powder per 1 gallon of water. Mix until dissolved. Administer at the indicated rates in the total water consumed over a full, 24-hour period. Administer 7 to 14 days.

NO COPY

Time/Date: NPVSOPMKN&CÑ=N=N=VLNVLNQ=UWNN=y

Project No.	Artwork Number	Description	Country
12017	13962301	Aureomycin	US
Dimensions		Drawing No.	SKU No.
10.053" x 18.25"		DWG-102290-02	3829000
Item		Foils	
Substrate: Aluminum		EDITOR'S COPY	
Colors: PMS 032 PMS 102 PMS 281 Printable Area		GS: DATE:	
Mgr	Rev	GA	PR
GS	2		
GA			
CHANGES OK		CHANGES OK	