

**INDICATIONS:**

For use in cattle as an aid in treating uncomplicated milk fever (parturient paresis).

**DOSAGE AND ADMINISTRATION:**

For intravenous administration only.

Cattle..... 250-500 mL  
Dosage may be repeated at 8 to 12 hour intervals if required.

**CAUTION:**

Solution should be warmed to body temperature and administered slowly. Entire contents should be used upon entering, as this product contains no preservative. Discard any unused portion. Do not use if solution is cloudy or contains precipitate.

**NOTE:**

Monitor animal's condition closely. Re-treatment may be necessary. If there is no noticeable improvement within 24 hours following treatment, consult your veterinarian. Aseptic precautions should be observed.

**Distributed by:**  
Aspen Veterinary Resources,\* Ltd.  
Liberty, MO 64068 USA

A140NT 01/15

**Manufactured by:**  
Nova-Tech, Inc.  
Grand Island, NE 68801 USA

18-802  
RMS# 92-503

NDC# 46066-802-50

**aspen**  
VETERINARY RESOURCES,\* LTD.



**Calcium Gluconate  
23% Solution**

**STERILE**



**For Animal Use Only  
Keep Out of Reach of Children**

**NET CONTENTS:  
500 mL (16.91 fl oz)**

**EACH 500 mL CONTAINS:**

Calcium Gluconate.....23% w/v  
(provides 10.71 g of Calcium\* per 500 mL)  
Water for Injection.....q.s.  
\* Present as salts or boryl esters of gluconic acid.

**Store between 15°C-30°C (59°F-86°F).**

TAKE TIME  OBSERVE LABEL DIRECTIONS



Lot No.

Exp. Date