

EACH 100 mL CONTAINS:

Active Ingredients

Calcium Gluconate
(provides 2.14 grams Calcium)23 grams

Inactive Ingredients

Water for Injectionq.s.
Contains boric acid as a solubilizing agent.

ELECTROLYTES PER 1000 mL:

Calcium 1069 mEq
Hypertonic

INDICATIONS: To be used as an aid in the treatment of parturient paresis (milk fever) in cattle.




MADE IN USA

Distributed by: MWI
Boise, ID 83705
www.VetOne.net

Manufactured by:
Nova-Tech, Inc.
Grand Island, NE 68801

RMS 92-1003
18-802
Rev. 05/23



3 13985 00900 6

NDC 13985-802-50

500 mL



Calcium Gluconate 23%

Solution, USP

**Sterile
Preservative Free**

For animal use only.

Keep out of reach
of children.



V1 510104

Net Contents: 500 mL (16.91 fl oz)

DOSAGE AND ADMINISTRATION: Cattle: 250 mL to 500 mL. Administer intravenously or intraperitoneally. Solution should be warmed to body temperature and administered slowly.

This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. Discard any unused portion. Handle under aseptic conditions. If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

For technical questions, call 1-833-833-4883.

STORAGE: Store between 15° and 30°C (59°–86°F).

TAKE TIME



OBSERVE LABEL
DIRECTIONS

LOT NO.:

EXP. DATE: