

**INDICATIONS:** To be used as an aid in the treatment of parturient paresis (milk fever) in cattle.

**DOSAGE AND ADMINISTRATION:**

Cattle: 250 to 500 mL

Administer intravenously, intraperitoneally or subcutaneously. If given subcutaneously, divide dosage among several injection sites and massage to aid in absorption.

Solution should be warmed to body temperature and administered slowly. This product is sterile in unopened container. It contains no preservative. Use entire contents when first opened. It should be handled under aseptic conditions.

If there is no apparent improvement in the animal's condition within 24 hours, consult a veterinarian.

**Store between 15°C and 30°C (59°F and 86°F).**

® Registered trademark of Agri Laboratories, Ltd.

18-802  
RMS 92-567

Rev. 03-14

**LOT**

**EXP**

NDC 57561-802-50

# CALCIUM GLUCONATE 23% SOLUTION

**Sterile Preservative Free**

**For Animal Use Only**

**Keep Out of Reach Of Children**

**NET CONTENTS: 500 mL**



**Each 100 mL CONTAINS:**

**ACTIVE INGREDIENTS**

Calcium Gluconate  
(provides 2.14 grams  
Calcium) ..... 23 grams  
Water for Injection ..... q.s.

**INACTIVE INGREDIENTS**

Boric Acid (as a  
solubilizing agent) ..... 3.5 grams  
Electrolytes per 1000 mL:  
Calcium 1069 mEq;  
Hypertonic ..... pH 2.5-4.5

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

**Manufactured for:**  
**Agri Laboratories, Ltd.**  
**St. Joseph, MO 64503**

