

NDC 54925-082-04

ANADA 200-420, Approved by FDA

CEFTIFLEX[®]

(ceftiofur sodium sterile powder)

4 grams

For intramuscular and subcutaneous injection in cattle only.
For intramuscular injection in swine, sheep, goats, and horses.

For subcutaneous injection only in day-old chickens,
and day-old turkey poults.

May Be Used in Lactating Dairy Cattle, Sheep, and Goats.

Caution: Federal (USA) law restricts this drug to use
by or on the order of a licensed veterinarian.
Reconstitute with 80 mL Sterile Water for Injection.
Shake thoroughly prior to use.

Each mL of the resulting solution contains ceftiofur
sodium equivalent to 50 mg ceftiofur.



For Once Daily Injection – See Package Insert

Horses	1.0 - 2.0 mg/lb	2 - 4 mL/100 lb
Day-old Chickens	0.08 - 0.2 mg/chick	1 mL/250 - 625 chicks
Sheep/Goats	0.5 - 1.0 mg/lb	1 mL/50 - 100 lb
Cattle	0.5 - 1.0 mg/lb	1 mL/50 - 100 lb
Swine	1.36 - 2.27 mg/lb	1 mL/22 - 37 lb
Day-old Poults	0.17 - 0.5 mg/poult	1 mL/100 - 294 poults

WARNING: Not for human use. Keep out of reach of children. To avoid possible allergic reactions, users are advised to avoid direct contact of this product with the skin or mucous membranes (see package insert).

Store **unreconstituted** product at controlled room temperature 20° to 25°C (68° to 77°F) [see USP]. Protect from light.

See package insert for complete product information and storage conditions.

Each vial contains: ceftiofur sodium equivalent to 4 grams ceftiofur.

RESIDUE WARNINGS: Pre-slaughter meat withdrawals: Cattle and Swine: 4 days after last treatment. Sheep, goats, day-old chickens and turkey poults: 0 hrs. Milk discard time: 0 hrs.



Mfd. for: Med-Pharmex Inc., Pomona, CA 91767
By: Cephalazone Pharma LLC, Pomona, CA 91767

Lot No./Exp. Date:

LA-0018