DIRECTIONS FOR USE

Dissolve the contents of one packet of Chlortetracycline Hydrochloride Soluble Powder in four (4) gallons of water to prepare a stock solution, when metered at the rate of 1 oz of stock solution per gallon of drinking water, will deliver 200 mg chlortetracycline hydrochloride per gallon of drinking water. One level measuring teaspoon contains approximately 360 mg chlortetracycline hydrochloride which will treat 36 lbs (16 kg) of animal at 10 mg/lb for one day. Not to be used for more than 14 consecutive days in chickens and turkeys and five days in swine. The concentration of drug required in drinking water must be adjusted to provide a correct dosage in order to compensate for variations in the age of the animal, class of poultry, environmental temperature and humidity; each of which affects water consumption.

CHICKENS AND TURKEYS

Administer for 7 to 14 days. Medicate continuously at the first clinical signs of disease. The dosage range permitted provides for different levels based on the severity of the infection. As a generalization, 100 turkeys will drink one gallon of water per day for each week of age. Chickens will consume one half this amount.

CONTENTS

This packet contains 102.4 grams of chlortetracycline hydrochloride (64 g/lb). Store at Controlled Room Temperature of 20° to 25°C (68° to 77°F). Brief exposure up to 40°C (104°F) may be tolerated.

TAKE TIME OBSERVE LABEL DIRECTIONS

100435

Lot No./Exp. Date

Rev. 08-16
5099-01

NDC 76475-236-01

CHLORTETRACYCLINE HYDROCHLORIDE SOLUBLE POWDER

Antibiotic

For Use in Drinking Water of Chickens, Turkeys and Swine.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

FOR VETERINARY USE ONLY

Restricted Drug(s) (California)
Not For Human Use

Store at Controlled Room Temperature of 20° to 25°C (68° to 77°F). Brief exposure up to 40°C (104°F) may be tolerated.

Prepare fresh drinking water solutions daily.

For use in drinking water only.

Not for use in liquid feed supplements, milk or milk replacers.

Use as the sole source of drinking water.

KEEP OUT OF REACH OF CHILDREN

Safety Data Sheet (SDS) available.

Net Weight: 725.8 g (25.6 oz)

ANADA 200-236, Approved by FDA

Manufactured For:
Quo Vademus, LLC
Kenansville, NC 28349
customercare@quovademusllc.com • 910-296-1632

SCAN FOR
SDS

QUOVADEMUS®

INDICATIONS FOR USE AND DOSAGE

CHICKENS

Control of infectious synovitis caused by *M. synoviae* - 200 to 400 mg chlortetracycline hydrochloride per gallon drinking water (5.0 - 13.4 mg/lb body weight per day). Control of chronic respiratory disease (CRD) and air sac infections caused by *M. gallisepticum* and *E. coli* - 400 to 800 mg chlortetracycline hydrochloride per gallon drinking water (10.0 - 26.8 mg/lb body weight per day).

TURKEYS

Control of infectious synovitis caused by *M. synoviae* - 400 mg chlortetracycline hydrochloride per gallon drinking water (3.2 - 16.8 mg/lb body weight per day). Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) - 25 mg chlortetracycline hydrochloride per pound body weight daily.

SWINE

Control and treatment of bacterial enteritis (scours) caused by *E. coli* and *Salmonella* spp. and bacterial pneumonia associated with *Pasteurella* spp., *Actinobacillus pleuropneumonia*, *Klebsiella* spp. - 10 mg chlortetracycline hydrochloride per pound body weight daily in divided doses for 3 to 5 days.

WARNING

Withdraw 24 hours before slaughter of chickens, turkeys and five days for swine. Do not feed to chickens producing eggs for human consumption. For growing turkeys only.

PRECAUTION

Prepare fresh drinking water solutions daily. In automatic watering systems, prepare fresh stock solutions every 6 hours. Solutions are not stable in galvanized containers; their use is not recommended. Drinking water solutions should not be stored at 40°C (104°F) for longer than 12 hours. Stock solutions should not be stored at 40°C (104°F) for longer than 3 hours. Use only against organisms sensitive to chlortetracycline. Use as the sole source of chlortetracycline. Do not mix in liquid feed supplements, milk or milk replacers. Administer 1 hour before or 2 hours after feeding with milk or milk replacers.

12.567" CUT OFF

1" | 2.375" | .25" | 6.125" | .25" | 2.375" | .5"

12.875" PACKAGE WIDTH

36.625" WEB WIDTH

MACHINE DIRECTION

LINE BLACK

PANTONE 364

DIELINE IS FOR LAYOUT PURPOSES ONLY!
DIELINE DOES NOT PRINT