

FOR ANIMAL USE ONLY**To prepare 50 Gallons of Medicated Water**

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of Cocci-Cure® P FOR POULTRY 9.6% Oral Solution in about 5 gallons of water in a 50 gallon medication barrel. Stir, then add water to the 50 gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use ½ pint (8 fluid ounces) of Cocci-Cure® P FOR POULTRY 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of Cocci-Cure® P FOR POULTRY 9.6% Oral Solution.

PREMEASURED FOR AUTOMATIC WATER PROPORTIONERS:

For use in automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

TO PREPARE DOSAGE LEVELS OF	Cocci-Cure® P FOR POULTRY 9.6% Oral Solution PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

Benzoic acid as Sodium Benzoate 0.1% added as preservative.

STORAGE: Store container in an upright position. Store between 5°C-25°C (41°F-77°F) with brief excursions to 40° C.

Cocci-Cure® is a registered trademark of Aspen Veterinary Resources, Ltd.

To obtain an MSDS or for assistance contact Aspen Veterinary Resources, ® Ltd. at 1-800-792-1238.



Cocci-Cure® P FOR POULTRY



**(amprolium)
9.6% Oral Solution
Coccidiostat**

Active Ingredient: amprolium.....9.6%

**Water-soluble treatment for coccidiosis
Treats 800 Gallons at 0.012% level**

**Restricted Drug (California) – Use Only as Directed
ANADA 200-496, Approved by FDA**

NET CONTENTS:

1 Gallon (128 fl oz) (3.785 L)

INDICATIONS: Cocci-Cure® P FOR POULTRY 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

USE DIRECTIONS: Give amprolium at the 0.012% level (8 fl oz Cocci-Cure® P FOR POULTRY 9.6% Oral Solution per 50 gallons) as soon as coccidiosis is diagnosed and continue for three to five days. (In severe outbreaks, give amprolium at the 0.024% level). Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time.

Use as the sole source of amprolium.

WARNING: Keep this and all drugs out of reach of children. NOT FOR HUMAN USE.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

Product No. 1AMP029
8AMP031 Rev: 03/15

Manufactured for:
Aspen Veterinary Resources,® Ltd.
Liberty, MO 64068
www.aspenveterinaryresources.com
A460B Rev. 03/15

Lot:
Exp:

