

FOR ANIMAL USE ONLY

INDICATIONS: In cattle, horses, swine and sheep as a topical antiseptic for use prior to surgical procedures such as castrating and docking; for ringworm and foot rot; for application to the navel of newborn animals; and for minor cuts, teat sores, bruises and abrasions.

DIRECTIONS: Remove spray cap from 16 oz. container and refill from gallon container. Replace spray cap tightly. Hold container approximately 4 to 6 inches from the area to be sprayed. Point valve at the area to be sprayed. Pull trigger of valve and spray area once lightly. May be repeated daily, if necessary, until the area is healed.

CAUTION: If redness, irritation or swelling persists or increases, discontinue use and consult a veterinarian. Not for use on burns or in body cavities or deep wounds.

NOTE: When used on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of milk.

TAKE TIME

OBSERVE LABEL
DIRECTIONS

Manufactured for:

DURVET, INC.

Blue Springs, Missouri 64014

Rev. 05-14

durvet**CONTROLLED
IODINE SPRAY**

Antiseptic/Disinfectant
For Skin Disinfection, Superficial Wounds, Navel Cords

CAUTION: KEEP OUT OF REACH OF CHILDREN**www.durvet.com****NET CONTENTS: 16 fl. oz. (473 mL)****KEEP OUT OF REACH
OF CHILDREN****INGREDIENTS:****Active Ingredients:**

Polyethoxylated nonylphenol iodine complex
(provides 1% titratable iodine) 5%

Inactive Ingredients: 95%
Nonionic Surfactant and Water.

CAUTION: Harmful or fatal if swallowed. Do not apply to the eyes, mucous membranes or large areas of abraded skin. Avoid inhalation of mist.

STORAGE: Store at controlled room temperature between 10°-30°C (50°-86°F). Keep tightly closed, protect from light.



Lot No.

Exp. Date