

PBOM 18x6x33w2.25"step

NADA 034-393
Approved by the FDA

COYDEN[®] 25%

(clopidol)

Type A Medicated Article

ACTIVE DRUG INGREDIENT: Clopidol.....113.5 gm/lb (25%)

INGREDIENTS: Rice Hulls, Mineral Oil, Fatty Acid Glycerides

BROILERS/REPLACEMENT CHICKENS:

As an aid in the prevention of cecal and intestinal coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. mivati*.

TURKEYS:

As an aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

SEE DIRECTIONS ON BACK PANEL

FOR THE MANUFACTURE OF COMPLETE FEEDS ONLY.

Manufactured by:
HUVEPHARMA, INC. St. Louis, MO 63116
Distributed by:
HUVEPHARMA, INC. Peachtree City, GA 30269
For questions, call 1-877-426-7765

Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269



® COYDEN is a registered trademark of Huvepharma, Inc.

Net Wt.: 50 lb (22.68 kg)



COYDEN[®] 25%
(clopidol)
Type A Medicated Article
HUVEPHARMA

LOT NO & EXP DATE

123456 7/09
FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
GUARANTEED BY
GATEWAY PACKAGING COMPANY

Restricted Drug (California)
Use only as Directed



INDICATIONS:

BROILER AND REPLACEMENT CHICKENS

As an aid in the prevention of coccidiosis in broilers and replacements for caged layers caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. mivati*.

TURKEYS GROWN FOR MEAT PURPOSES ONLY

As an aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

WARNING:

Keep this and all drugs out of the reach of children.

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, obtain an MSDS or for assistance, contact Huvepharma at 1-877-426-7765.

RESIDUE INFORMATION:

CHICKENS: When feeding 0.025% clopidol, reduce dosage to 0.0125% or withdraw medication 5 days before slaughter. Do not feed to chickens over 16 weeks of age.

TURKEYS: Withdraw medication 5 days before slaughter.

CAUTION: Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period.

MIXING DIRECTIONS:

For Complete Rations - Type C Medicated Products
COYDEN[®] 25% (clopidol) should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed. COYDEN 25% may be used to manufacture a Type C medicated feed in the concentration range of 0.0125% to 0.025% clopidol.

To aid in even distribution of COYDEN 25% in the finished feed, prepare a mixture of COYDEN 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and mix thoroughly.

DIRECTIONS FOR USE:

Must be mixed with other ingredients before feeding.

Mixing and conveying equipment should be properly grounded.

Use 1 pound of COYDEN 25% per ton of feed to produce a feed containing 0.0125% clopidol.

Use 2 pounds of COYDEN 25% per ton of feed to produce a feed containing 0.025% clopidol.

FEEDING DIRECTIONS

Feed continuously, as the sole ration, either the 0.0125% or 0.025% Type C medicated feed depending upon management practices, degree of exposure, and amount of feed eaten.

STORE IN A COOL PLACE
KEEP BAG TIGHTLY CLOSED

COYDEN[®] 25%
(clopidol)
Type A Medicated Article

5031-01
Rev. 07-2009