

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND  
DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

<b>FIRST AID</b>	
<b>If in eyes</b>	-Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.
<b>If swallowed</b>	-Call poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to do so by the poison control center or doctor. -Do not give anything by mouth to an unconscious person.
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-292-8680 for emergency medical treatment information.	

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Keep container tightly closed when not in use. Store in original container, in a dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** This product is contained in a nonrefillable container. Do Not reuse or refill this container.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See side panel for Application Instructions. Personal Protection Equipment (PPE) is to be worn by all applicators and handlers of this product. PPE shall include protective gloves and either a dust mask or dust/mist filtering respirator.

**Darkling Beetle Control  
An Insecticide for the Control  
of Darkling Beetles, Hide  
Beetles, and Flies.**

**Active Ingredient:**

Boric Acid .....100%  
CAS# 10043-35-3

**KEEP OUT OF THE REACH  
OF CHILDREN  
CAUTION**

See Side Panel for Additional  
Precautionary Statements

EPA Reg No. 67546-2  
EPA Est. 67542-CHL-001

**Manufactured by:  
National BoraXX Corp.  
3690 Orange Place, Suite 480  
Cleveland, Ohio 44122  
Emergency Phone: 1-800-292-8680**

®Reg. Trademark of National BoraXX Corp.  
©1995 National BoraXX Corp.

**Net Contents: 50 lbs. (22.7 Kg.)**

**APPLICATION INSTRUCTIONS**

FOR DARKLING AND LESSER MEALWORM BEETLE, HIDE BEETLE, AND FLIES (Adults and Larvae).

For Poultry houses/hogbarns/barns in which birds/hogs are grown:

**DRY APPLICATION**

Before application of the Darkling Beetle Control, **remove animals.** Apply this product uniformly to floors of poultry house/hogbarns/barns or to old litter. Use a fertilizer or seed spreader to apply this product at the rate of 1-2 pounds for each 100 square feet, in bands, along feeder lines. Spread fresh litter of at least four inches depth uniformly over all treated areas (floor or old litter). Then introduce animals. Reapply after each grow-out, if needed, in the order stated above.

**WET APPLICATION**

Prior to application of Darkling Beetle Control, **remove animals from poultry house/hogbarns/barns.** Mix Darkling Beetle Control at the rate of 1 pound per 3 gallons of water. Apply the 3 gallons of mix for each 100 square feet of treated surface. Using appropriate low pressure spray equipment, saturate side walls, top plates, posts, and framing. Spray mix into cracks and crevices around insulation. Do not expose animals to spray. Reapply after each grow-out, if needed, in the order of steps outlined above. However, if the facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied. Darkling Beetle Control can be applied directly to manure for fly control.

Treatment of poultry houses in which birds are grown in cages (layer or high rise 'pit type' houses) should follow the following instructions.

**DRY APPLICATION**

Prior to application of the Darkling Beetle Control, **remove birds.** Apply at the rate of 1-2 pounds of Darkling Beetle Control for each 100 square feet of treated surface. Using low pressure hand wands, thoroughly dust side walls, top plates, posts, and framing. Treat into cracks and crevices around insulation. Then introduce birds. Reapply after each grow-out, if needed, in the order of steps outlined above. However, if the facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied to maintain Darkling Beetle control. Darkling Beetle Control can be applied directly to manure for fly control.

**NOTICE**

Manufacturer shall not be liable in respect to any injury or consequential damages suffered based on the use of this product contrary to the instructions given on this label.  
8/31/2016