

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Keep away from children, domestic animals and pets. Causes moderate eye irritation. Do not get in eyes or on clothing. All handlers (including applicators) must wear long pants, shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Users should remove clothing/ PPE immediately if pesticide gets inside. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may also be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

READ THIS LABEL: READ THIS ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH USE DIRECTIONS, USE RESTRICTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USE ONLY FOR THE SITES, PESTS, AND APPLICATION METHODS DESCRIBED ON THIS LABEL.

It is illegal to sell these blocks individually.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent exposure:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, nontarget mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around agricultural buildings and manmade agricultural structures: House mouse (*Mus musculus*), Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*), Cotton rat* (*Sigmodon hispidus*), Eastern harvest mouse* (*Reithrodontomys humuli*), Golden mouse* (*Ochrotomys nuttalli*), Polynesian rat* (*Rattus exulans*), Meadow vole* (*Microtus pennsylvanicus*), White-throated woodrat* (*Neotoma albigula*), Southern plains woodrat* (*Neotoma micropus*), and Mexican woodrat* (*Neotoma mexicana*). This product must be used in and within 100 feet of agricultural buildings and manmade agricultural structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Fence and perimeter baiting beyond 100 feet from an agricultural structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with DeciMax Blocks is prohibited.

DeciMax[®] Blocks

**KILLS RATS, HOUSE MICE,
AND MEADOW VOLES***

**For Indoor and Outdoor Rat, House
Mouse, and Meadow Vole* Control**

*This product may only be used inside and within 100 feet of
agricultural buildings and manmade agricultural structures*

**NORWAY RATS, ROOF RATS AND HOUSE MICE MAY CONSUME A LETHAL
DOSE IN A SINGLE NIGHT'S FEEDING WITH FIRST DEAD RATS AND MICE
APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS.**

ACTIVE INGREDIENT:

Bromadiolone 0.005%

OTHER INGREDIENTS:** 99.995%

TOTAL: 100.000%

**Contains denatonium benzoate

**This product has been treated with S-methoprene to protect the bait from infestation by insects.

EPA Reg. No. 270-373-61282 EPA Est. No. 61282-WI-01

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See side panel for additional Precautionary Statements

FIRST AID

Have label with you when obtaining treatment advice.

If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-498-5743** for emergency medical treatment information.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K, intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or **1-800-498-5743** at once.

NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**Net Contents: 9 lbs. (144 X 1 oz. blocks)
Net Weight: 9 lbs.**



(Directions for Use, continued)

Do not use in homes or other human residences. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. Do not broadcast bait. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 8 pounds of bait.

SELECTION OF TREATMENT AREAS

Determine areas where rats, mice, or meadow voles* will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rats, mice, or meadow voles* or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

Each DeciMax Block weighs 1 oz.

RATS: Place 3 to 16 bait blocks (at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE AND MEADOW VOLES*: Place one block per placement. Space placements at 8 to 12 foot intervals in infested areas. Two blocks may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole* activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all exposed dead animals and leftover bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed.

CONDITIONS OF SALE

To the extent consistent with applicable law, Seller warrants this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions as specified on the label. There are no other warranties, expressed or implied, concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

For non-emergency (e.g., current product information,
Call: **1-800-621-8829**

Manufactured by:

HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956 U.S.A.
HACCO, Inc. is a wholly-owned subsidiary of Neogen Corporation.

LA220512-0318

