



Denagard™ 10

(tiamulin hydrogen fumarate)

Type B Medicated Swine Feed

For manufacture of Type C medicated swine feed only

ACTIVE INGREDIENT: Tiamulin hydrogen fumarate: 10 g/lb

Ingredients: rice hulls, mineral oil, porcine gelatin

See reverse side for use directions, cautions and warnings.

Net Weight: 35 lb (15.9 kg)

Do not feed undiluted.

| Indications | Amount of Type B Medicated Feed Per Ton of Feed | Tiamulin hydrogen fumarate in Complete Type C Medicated Feed Per Ton | Withdrawal Period (days) |
|--|---|--|--------------------------|
| For treatment of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> susceptible to Tiamulin | 20 lb | 200 g | 7 |
| For control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellularis</i> | 3.5 lb | 35 g | 2 |
| For control of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> susceptible to Tiamulin | 3.5 lb | 35 g | 2 |

Warnings:

▶ Observe withdrawal time shown in table above. ◀

Keep out of reach of children. Avoid contact with skin. Direct contact with skin or mucous membranes may cause irritation.

Caution:

Do not feed undiluted. Do not use in feeds for animals other than swine. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Swine being treated with Denagard (tiamulin) should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use.

MIXING DIRECTIONS

Mix this medicated feed (Type B) with non-medicated feed ingredients to make a Type C medicated feed as follows:

| Type C Medicated Feed Tiamulin grams/ton Feed | Pounds Medicated Feed (Type B) | Pounds Non-medicated Feed* |
|---|--------------------------------|----------------------------|
| 35 | 3.5 | 1996.5 |
| 200 | 20 | 1980 |

*When making an approved Type C medicated feed drug combination, the inclusion rate of this non-medicated feed should be adjusted for the amount of the other drug added to make such combination Type C medicated feed.

Storage:

Store at or below 25°C (77°F). Temporary excursions to 40°C (104°F) permitted.

To report adverse effects, access medical information, or obtain additional product information call 1-800-428-4441

For a complete listing of adverse reactions for DENAGARD reported to CVM see: <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Elanco, Denagard and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Restricted Drug (California): Use only as directed.

NADA 139-472, Approved by FDA.

BG100138AMX

Marketed by:
Elanco Animal Health,
Indianapolis, IN 46285

