

For intravenous or intramuscular injection.

**USUAL DOSE:**

Bovine - 5 to 20 mg, Equine - 2.5 to 5 mg

**Warning:** A withdrawal period has not yet been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Read accompanying directions carefully.

**FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**

Dexium is a Trademark of Bimeda, Inc.

Manufactured by: Bimeda-MTC Animal Health Inc.

Cambridge, Ontario, Canada N3C 2N4

Bimeda, Inc. and Bimeda-MTC Animal Health Inc.  
are Divisions of Cross Vetpharm Group Ltd.

# Dexium™

## DEXAMETHASONE SOLUTION

**Veterinary 2 mg per mL**

**Sterile**

**CAUTION:** Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA #200-312 Approved by FDA

**Net Contents: 100 mL**

Manufactured for:  
Bimeda, Inc.  
Le Sueur, MN 56058



**Bimeda**

Each mL contains: 2 mg dexamethasone, 500 mg polyethylene glycol 400, 9 mg benzyl alcohol, 1.8 mg methylparaben and 0.2 mg propylparaben as preservatives, 4.75% alcohol, HCl to adjust pH to approximately 4.9, water for injection q.s.

**Store between 2° and 30° C (36° and 86°F).**

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Product No. 1DEX022

8DEX018D

UPC 80%  
8 99855 60015 8