



NDC 23243-370-07

# DI-METHOX<sup>®</sup> (sulfadimethoxine) CONCENTRATED SOLUTION 12.5%

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Antibacterial for use in drinking water  
For Oral Use in Chickens,  
Turkeys and Cattle

3.75 g sulfadimethoxine/fl oz  
Protect From Light

Approved by FDA under ANADA # 200-030

**NET CONTENTS: 3.8 Liters (1 GALLON)**



#### INDICATIONS:

**BROILER AND REPLACEMENT CHICKENS**– Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

**MEAT PRODUCING TURKEYS**– Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

**DAIRY CALVES, DAIRY HEIFERS AND BEEF CATTLE**– Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Fusobacterium necrophorum* (*Sphaerophorus necrophorus*) sensitive to sulfadimethoxine.

**PRECAUTION:** Store at room temperature; if freezing occurs, thaw before using. Protect from light; direct sunlight may cause discoloration. Freezing or discoloration does not affect potency. Prepare a fresh stock solution daily.

**CHICKENS AND TURKEYS**– If animals show no improvement within 5 days, discontinue treatment and re-evaluate diagnosis. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication. Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

**CATTLE**– During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. Treatment should not be continued beyond 5 days.

#### RESIDUE WARNINGS:

**CHICKENS AND TURKEYS**– Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

**CATTLE**– Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

STORE AT CONTROLLED ROOM TEMPERATURE 20-25°C (68-77°F).

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) – Use Only as Directed

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

5-3369-07  
Rev. 07-2021

Manufactured for  
Huvepharma, Inc.  
Peachtree City, GA 30269 <sup>®</sup> Registered trademark of Huvepharma, Inc.

LOT NO.:

EXP. DATE:



#### Dosage and Administration

##### Chickens and Turkeys

**Treatment Period:** 6 consecutive days

**Recommended Concentration:** Chickens–0.05%; Turkeys–0.025%

**Chickens** Add 1 fl oz (30 mL) to 2 gal of drinking water or 25 fl oz to 50 gal of drinking water.

**Turkeys** Add 1 fl oz (30 mL) to 4 gal of drinking water or 25 fl oz to 100 gal of drinking water.

**Automatic Proportioners\* Stock Solution**–To make 2 gal of stock solution:

**Chickens** Add 1 gal of DiMethox<sup>®</sup> Concentrated Solution 12.5% to 1 gal of water.

**Turkeys** Add 2 qt of DiMethox<sup>®</sup> Concentrated Solution 12.5% to 6 qt of water.

\*Set proportioner to a feed rate of 1 fl oz of stock solution per gal of water

##### Dairy Calves, Dairy Heifers, and Beef Cattle

**Treatment Period:** 5 consecutive days

**Dosage:** Initial dose of 25 mg/lb followed by four maintenance doses of 12.5 mg/lb/day

##### Summer Administration

Dosage recommendations for summer are based on an estimated water intake of 1 gal of water for every 100 lb of body weight per day.

Daily Drinking Water Supply	DiMethox <sup>®</sup> Concentrated Solution 12.5%	
	Initial Dose	Maintenance Dose
25 gal	1 pt (16 fl oz)	1 cup (8 fl oz)
50 gal	1 qt (32 fl oz)	1 pt (16 fl oz)
200 gal	1 gal (128 fl oz)	2 qt (64 fl oz)

##### Winter Administration

Dosage recommendations for winter are based on an estimated water intake of 1 gal of water for every 150 lb of body weight per day.

Daily Drinking Water Supply	DiMethox <sup>®</sup> Concentrated Solution 12.5%	
	Initial Dose	Maintenance Dose
16 gal	1 pt (16 fl oz)	1 cup (8 fl oz)
33 gal	1 qt (32 fl oz)	1 pt (16 fl oz)
127 gal	1 gal (128 fl oz)	2 qt (64 fl oz)

For individual treatment of cattle, Di-Methox<sup>®</sup> Concentrated Solution 12.5% may be given as a drench.

Administer using an initial dose of 25 mg/lb followed by 4 maintenance doses of 12.5 mg/lb/day. One fl oz will medicate one 150-lb animal initially and 1/2 fl oz will medicate one 150-lb animal on maintenance dose.



3 57561 00077 3