

INDICATIONS:

BROILER AND REPLACEMENT CHICKENS- Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

MEAT PRODUCING TURKEYS- Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

DAIRY CALVES, DAIRY HEIFERS AND BEEF CATTLE- Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella spp.* sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Fusobacterium necrophorum (Sphaerophorus necrophorus)* sensitive to sulfadimethoxine.

PRECAUTION: Store at room temperature; if freezing occurs, thaw before using. Protect from light; direct sunlight may cause discoloration. Freezing or discoloration does not affect potency. Prepare a fresh stock solution daily.

CHICKENS AND TURKEYS- If animals show no improvement within 5 days, discontinue treatment and re-evaluate diagnosis. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication.

Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

CATTLE- During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. Treatment should not be continued beyond 5 days.

RESIDUE WARNINGS:

CHICKENS AND TURKEYS- Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

CATTLE- Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.

STORE AT CONTROLLED ROOM TEMPERATURE 15°-30°C (59°-86°F).

TAKE TIME



OBSERVE LABEL DIRECTIONS

FOR USE IN ANIMALS ONLY

NOT FOR HUMAN USE

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) – Use Only as Directed

S-3369-07
Rev. 07-16

®Registered trademark of Agri Laboratories, Ltd.

Manufactured for:
Agri Laboratories, Ltd.
St. Joseph, MO 64503

LOT NO.:

EXP. DATE:

NDC 57561-370-07

DI-METHOX[®]**(sulfadimethoxine)****CONCENTRATED SOLUTION 12.5%**

CAUTION: Federal (USA) law restricts
this drug to use by or on the order
of a licensed veterinarian.

Antibacterial for use in drinking water

**For Oral Use in Chickens,
Turkeys and Cattle**

3.75 g sulfadimethoxine/fl oz

Protect From Light

ANADA 200-030, Approved by FDA

NET CONTENTS: 3.8 Liters (1 GALLON)**AgriLabs[®]**Making A Healthy Difference[®]**Dosage and Administration****Chickens and Turkeys****Treatment Period: 6 consecutive days****Recommended Concentration: Chickens–0.05%; Turkeys–0.025%**Chickens Add 1 fl oz (30 mL) to 2 gal of drinking water or
25 fl oz to 50 gal of drinking water.Turkeys Add 1 fl oz (30 mL) to 4 gal of drinking water or
25 fl oz to 100 gal of drinking water.**Automatic Proportioners*Stock Solution–To make 2 gal of stock solution:**Chickens Add 1 gal of DiMethox[®] Concentrated Solution 12.5% to 1 gal of water.Turkeys Add 2 qt of DiMethox[®] Concentrated Solution 12.5% to 6 qt of water.

*Set proportioner to a feed rate of 1 fl oz of stock solution per gal of water

Dairy Calves, Dairy Heifers, and Beef Cattle**Treatment Period: 5 consecutive days**

Dosage: Initial dose of 25 mg/lb followed by four maintenance doses of 12.5 mg/lb/day

Summer Administration

Dosage recommendations for summer are based on an estimated water intake of 1 gal of water for every 100 lb of body weight per day.

DiMethox[®] Concentrated Solution 12.5%

| Daily Drinking Water Supply | Initial Dose | Maintenance Dose |
|-----------------------------|-------------------|------------------|
| 25 gal | 1 pt (16 fl oz) | 1 cup (8 fl oz) |
| 50 gal | 1 qt (32 fl oz) | 1 pt (16 fl oz) |
| 200 gal | 1 gal (128 fl oz) | 2 qt (64 fl oz) |

Winter Administration

Dosage recommendations for winter are based on an estimated water intake of 1 gal of water for every 150 lb of body weight per day.

DiMethox[®] Concentrated Solution 12.5%

| Daily Drinking Water Supply | Initial Dose | Maintenance Dose |
|-----------------------------|-------------------|------------------|
| 16 gal | 1 pt (16 fl oz) | 1 cup (8 fl oz) |
| 33 gal | 1 qt (32 fl oz) | 1 pt (16 fl oz) |
| 127 gal | 1 gal (128 fl oz) | 2 qt (64 fl oz) |

For individual treatment of cattle, Di-Methox[®] Concentrated Solution 12.5% may be given as a drench.

Administer using an initial dose of 25 mg/lb followed by 4 maintenance doses of 12.5 mg/lb/day. One fl oz will medicate one 150-lb animal initially and ½ fl oz will medicate one 150-lb animal on maintenance dose.



3 57561 00077 3