

14081001  
LOT/EXP

100 ML  
**EXCENEL® RTU EZ**  
**Ceftiofur Hydrochloride**  
Sterile Suspension

Equivalent to  
**50MG PER ML CEFTIOFUR**

For intramuscular injection in swine.

For intramuscular and subcutaneous injection in cattle.  
This product may be used in lactating dairy cattle.

Not for use in calves to be processed for veal.

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits extra-label use of this drug in cattle and swine for disease prevention purposes; at unapproved doses, frequencies, durations, or routes of administration; and in unapproved major food producing species/production classes.

For Use in Animals Only

NADA 141-285, Approved by FDA



zoetis

15280000

For Once Daily Injection — See Package Insert

1 mL/90 - 100 lb	0.5 - 1 mg/lb	Cattle
1 mL/2.2 - 37 lb	1.36 - 2.27 mg/lb	Swine

Warning: Not for human use. Keep out of reach of children. To avoid possible allergic reactions, users are advised to avoid direct contact of this product with the skin or other mucous membranes. See package insert for complete product information.

**RESIDUE WARNINGS**

**Swine:** When used according to label indications, dosage and route of administration, treated swine must not be slaughtered for 4 days following the last treatment.

Use of dosages in excess of those indicated or by unapproved routes of administration may result in illegal residues in edible tissues. **Cattle:** When used according to label indications, dosage and route of administration, treated cattle must not be slaughtered for 4 days following the last treatment. When used according to label indications, dosage and route of administration, a milk discard time is not required. Uses of dosages in excess of those indicated or by unapproved routes of administration, such as intramammary, may result in illegal residues in edible tissues and/or milk. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

Store at controlled room temperature 20° to 25° C (68° to 77° F); excursions permitted 15° to 40° C (59° to 104° F). Protect from freezing. Shake well before using. Contents should be used within 42 days after the first dose is removed.

Each mL contains: ceftiofur hydrochloride equivalent to 50 mg ceftiofur, 2.50 mg polyoxyethylene sorbitan monooleate (polyorbate 80), 6.5 mg water for injection in a caprylic/capric triglyceride (Niglyox® 812) suspension.

Distributed by: Zoetis Inc., Kalamazoo, MI 49007