

RESIDUE WARNINGS: Cattle must not be slaughtered for human consumption within 4 days of the last treatment. Milk that has been taken during treatment and for 36 hours after the last treatment must not be used for food. Not for use in dry dairy cows. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal. Not for use in horses intended for food. Approved only for intravenous administration in cattle. Intramuscular administration has resulted in violative residues in the edible tissues of cattle sent to slaughter.

 **Bimeda**

Flunazine[®]

(flunixin meglumine)

INJECTABLE SOLUTION

50 mg/mL

Sterile Solution

Veterinary

Multi-Dose Vial

NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA# 200-387, Approved by FDA



Net
Contents: 100 mL

STORE BETWEEN 2°C - 30°C (36°F - 86°F).
PROTECT FROM FREEZING.

For Intravenous or Intramuscular Use in Horses and for Intravenous Use in Beef and Dairy Cattle. Not for Use in Dry Dairy Cows and Veal Calves.

Read accompanying directions carefully.

Flunazine[®] is a Registered Trademark of Bimeda, Inc.

Manufactured by:
Bimeda-MTC Animal Health Inc.
Cambridge, ON Canada

Manufactured for:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

Product No. 1FLU003 8FLU006D Rev. 05/12



 **PULL**

TAKE TIME

OBSERVE LABEL DIRECTIONS