

Anti-infective for day-old chickens and 1- to 3-day-old turkeys.

Indications: Chickens: GARASOL Injection is recommended for the prevention of early mortality in day-old chickens associated with *Escherichia coli*, *Salmonella typhimurium*, and *Pseudomonas aeruginosa* susceptible to gentamicin sulfate.

Turkeys: as an aid in the prevention of early mortality of 1- to 3-day-old turkeys associated with *Arizona paracolon* infections susceptible to gentamicin sulfate.

Dose and Administration: Chickens: Inject each day-old chicken subcutaneously in the neck with GARASOL Injection diluted with sterile physiologic saline solution to provide 0.2 mg gentamicin in a 0.2-mL dose. Turkeys: Inject each 1- to 3-day-old turkey subcutaneously in the neck with GARASOL Injection diluted with sterile physiologic saline solution to provide 1 mg gentamicin in a 0.2-mL dose.

Residue Warnings: For use in day-old chickens only.

Injected chickens must not be slaughtered for food for at least five (5) weeks following treatment. For use in 1- to 3-day-old turkeys only. Injected turkeys must not be slaughtered for food for at least nine (9) weeks following treatment. Store between 2° and 30°C (36° and 86°F).



100 mg/mL

Sterile

NDC 0061-0134-01

Garasol®

(GENTAMICIN SULFATE, USP)

Injection

NADA #101-862,
Approved by FDA.

(100 mL)

Each mL contains: gentamicin sulfate, USP equivalent to 100 mg gentamicin base; 3.2 mg sodium bisulfite; 0.1 mg edetate disodium; 4.5 mg sodium acetate, anhydrous; 3.0 mg glacial acetic acid; 0.8 mg methylparaben and 0.1 mg propylparaben as preservatives; water for injection q.s.

**Read Product Information sheet carefully.
FOR SUBCUTANEOUS USE ONLY.**

Made in Germany.

Rev 8/14

Copyright © 1987, 2006,
Intervet Inc. (d/b/a Merck Animal Health)
Madison, NJ 07940
a subsidiary of Merck & Co.
All rights reserved.

**Restricted Drug (California) –
use only as directed.**

LOT:
EXP:

Varnish and
Ink free

