

For use in swine drinking water only.

INDICATIONS: GEN-GARD Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSE: Colibacillosis - 1 level scoop per 240 gallons of drinking water for 3 consecutive days, 0.5 mg/lb/day. For proportioner use, add 1 level scoop to 2 gallons stock solution and dispense at the rate of 1 oz./gal. drinking water (25 mg/gal).

Swine Dysentery - 1 level scoop per 120 gallons of drinking water for 3 consecutive days, 1.0 mg/lb/day. For proportioner use, add 1 level scoop to 1 gallon stock solution and dispense at the rate of 1 oz./gal. drinking water (50 mg/gal).

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**



G-7473-03

Rev. 04-2022



NDC 23243-4740-3

GEN-GARD[®]
(gentamicin sulfate)
Soluble Powder
Antibacterial

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**Contains 120 grams of
Gentamicin per jar**

Approved by FDA under ANADA # 200-185

**NET WEIGHT:
360 g (12.7 oz.)**



Each gram contains: gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: Swine treated with the recommended dose of GEN-GARD Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

Read package insert directions carefully. Store between 2° and 30°C (36° and 86°F). For use in swine drinking water only. Restricted drug (California)- Use only as directed.

© Registered trademark of Huvepharma, Inc.
Manufactured for
Huvepharma, Inc.
Peachtree City, GA 30269

Lot No.

Exp. Date

GEN-GARD[®]

(gentamicin sulfate)

Soluble Powder

Antibacterial

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For Swine Drinking Water Only.

DESCRIPTION: Each gram of GEN-GARD Soluble Powder contains gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: Swine treated with the recommended dose of GEN-GARD Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

INDICATIONS: GEN-GARD Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSAGE AND ADMINISTRATION: GEN-GARD Soluble Powder is available in a 360 gram jar. The 360 gram jar contains a scoop that provides approximately 18 grams of product when level full.

For Control and Treatment of Colibacillosis - Administer GEN-GARD Soluble Powder in drinking water at the recommended level of 25 mg per gallon (1 scoop/240 gallons) for three consecutive days.

For Control and Treatment of Swine Dysentery - Administer GEN-GARD Soluble Powder in drinking water at the recommended level of 50 mg per gallon (1 scoop/120 gallons) for three consecutive days.

The concentration of the medication should be adjusted in extremely hot or cold weather to ensure a gentamicin dosage of approximately 0.5 mg/lb/day for 3 days for colibacillosis or 1.0 mg/lb/day for 3 days for swine dysentery. A 50-lb pig will, under average conditions, consume 0.9 gallon of water per day.

For swine dysentery, if the condition recurs, the medication may be repeated. Because of the tendency for the disease to recur on a premise with a history of swine dysentery, a control program should be used following treatment.

DIRECTIONS FOR USE: To prepare medicated water, mix GEN-GARD Soluble Powder and drinking water according to the following tables:

Proportioner Use:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GEN-GARD to 2 gallons of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 25 mg/gallon.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GEN-GARD to 1 gallon of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 50 mg/gallon.

Bulk Preparation:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GEN-GARD to 240 gallons of water.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GEN-GARD to 120 gallons of water.

Medicated drinking water should not be stored or offered in rusty containers since the drug is quickly destroyed in such containers.

Medicated water should be prepared daily and be the sole source of drinking water for 3 consecutive days.

The daily water consumption figures shown below are approximations for mild temperatures and are presented as a guide only. Actual water consumption varies with environmental temperature, humidity, and diet.

Approximate Daily Water Consumption

<u>Weight of Pig (lb)</u>	<u>Gallons for 5 Pigs</u>	<u>Gallons for 10 Pigs</u>
25	3.0	6.0
50	4.5	9.0
75	6.0	12.0
100	7.5	15.0

SIDE EFFECTS: There have been no reported side effects following the recommended use of GEN-GARD Soluble Powder in swine.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

CONTRAINDICATIONS: There are no known contraindications to this drug when used as directed.

STORAGE CONDITIONS: Store between 2° and 30°C (36° and 86°F)

HOW SUPPLIED: GEN-GARD Soluble Powder is available in a 360 gram jar with 2 jars per shipper.

Approved by FDA under ANADA # 200-185

**Not for Use in Humans
Keep Out of Reach of Children
Restricted Drug (California) -
use only as directed**

Manufactured for
Huvepharma, Inc.
Peachtree City, GA 30269

Rev. 04-2022