


**INDICATIONS:** GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

**DOSE:**

*Colibacillosis* – 1 level scoop per 240 gallons of drinking water for 3 consecutive days, 0.5 mg/lb/day. For proportioner use, add 1 level scoop to 2 gallons stock solution and dispense at the rate of 1 oz/gal drinking water (25 mg/gal).

*Swine Dysentery* – 1 level scoop per 120 gallons of drinking water for 3 consecutive days, 1.0 mg/lb/day. For proportioner use, add 1 level scoop to 1 gallon stock solution and dispense at the rate of 1 oz/gal drinking water (50 mg/gal).

GentaMed is a Trademark of Bimeda, Inc.

 Bimeda

# GentaMed™ Soluble Powder

(Gentamicin Sulfate)

Contains 120 grams  
of Gentamicin per jar

**Antibacterial**

**FOR ANIMAL USE ONLY  
KEEP OUT OF REACH  
OF CHILDREN  
Restricted Drug (California)  
- Use Only as Directed**

**CAUTION: Federal law  
restricts this drug to use  
by or on the order of a  
licensed veterinarian.**

ANADA 200-494, Approved by FDA



**Net Weight:  
360 g (12.7 oz)**

Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058  
www.bimeda.com

**EACH g CONTAINS:** Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

**RESIDUE WARNING:** Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

Read accompanying directions carefully.  
Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

For use in swine drinking water only.  
8GEN001 Rev: 05/16 Product No. 1GEN009



**PULL HERE  
TO OPEN**

**TAKE TIME  
  
OBSERVE LABEL  
DIRECTIONS**

LOT:

EXP: