

Gil 1% Iodine Wound Spray

Topical Antiseptic

For Animal Use Only

**WARNING Causes Eye Irritation
May Be Harmful If Swallowed**

Keep Out of Reach of Children

For use on horses, cattle, swine and sheep as a topical antiseptic prior to surgical procedure such as castrating and tail docking; for application to the umbilical stump (naval) of newborn animals; and for the aid in treatment of minor cuts, bruises, teat sores, and abrasions.

Active Ingredient

NPE-free

Iodine* 1.00%
From alkyl alcohol ethoxylated iodine complex (as iodophor)

Inactive Ingredients

Water, nonionic surfactant, pH buffering agent 99.00%

FIRST AID

EYES: Immediately flush eyes with plenty of water for at least 15 minutes occasionally lifting the upper and lower lids. Contact a physician immediately.

INGESTION: Do NOT induce vomiting. Drink large amounts of milk or water. Contact a physician immediately. Do not give anything by mouth to an unconscious person.



Gilmer Industries, Inc.

560 Stone Spring Road PO Box 1247
Harrisonburg, VA 22801 Harrisonburg, VA 22803
Phone 540/434-8877

Precautions

- KEEP FROM FREEZING. If frozen, discard.
- Do NOT get in eyes or on clothing. Do NOT take internally.
- Do NOT allow to contaminate food. Wash thoroughly after handling.
- Keep container tightly closed when not in use.
- Store in a cool, dry, well-ventilated area out of direct sunlight.

Caution

- If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.
- Not for use on burns or in body cavities or deep wounds.
- If used on or near the teats or udders of dairy animals, wash teats and udders thoroughly before milking to prevent milk contamination.

Directions

Fill dedicated sprayer with product. Hold approximately 4-6 inches from area to be sprayed.

- For abraded skin, spray area once lightly. Repeat daily as necessary until area has healed.
- For the umbilical stump of a newborn animal, apply 2-3 times a day until the stump has dried.
- For use prior to a surgical procedure, allow a 10-minute contact time.

Do NOT return unused product to original container.

Net Content: **ONE Gallon (3.78L)**

K20Lot: -2020

Expiration: -2022

Gil 1% Iodine Wound Spray

Topical Antiseptic

For Animal Use Only

**WARNING Causes Eye Irritation
May Be Harmful If Swallowed**

Keep Out of Reach of Children

For use on horses, cattle, swine and sheep as a topical antiseptic prior to surgical procedure such as castrating and tail docking; for application to the umbilical stump (naval) of newborn animals; and for the aid in treatment of minor cuts, bruises, teat sores, and abrasions.

Active Ingredient

NPE-free

Iodine* 1.00%
From alkyl alcohol ethoxylated iodine complex (as iodophor)

Inactive Ingredients

Water, nonionic surfactant, pH buffering agent 99.00%

FIRST AID

EYES: Immediately flush eyes with plenty of water for at least 15 minutes occasionally lifting the upper and lower lids. Contact a physician immediately.

INGESTION: Do NOT induce vomiting. Drink large amounts of milk or water. Contact a physician immediately. Do not give anything by mouth to an unconscious person.



Gilmer Industries, Inc.

560 Stone Spring Road PO Box 1247
Harrisonburg, VA 22801 Harrisonburg, VA 22803
Phone 540/434-8877

Precautions

- KEEP FROM FREEZING. If frozen, discard.
- Do NOT get in eyes or on clothing. Do NOT take internally.
- Do NOT allow to contaminate food. Wash thoroughly after handling.
- Keep container tightly closed when not in use.
- Store in a cool, dry, well-ventilated area out of direct sunlight.

Caution

- If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.
- Not for use on burns or in body cavities or deep wounds.
- If used on or near the teats or udders of dairy animals, wash teats and udders thoroughly before milking to prevent milk contamination.

Directions

Fill dedicated sprayer with product. Hold approximately 4-6 inches from area to be sprayed.

- For abraded skin, spray area once lightly. Repeat daily as necessary until area has healed.
- For the umbilical stump of a newborn animal, apply 2-3 times a day until the stump has dried.
- For use prior to a surgical procedure, allow a 10-minute contact time.

Do NOT return unused product to original container.

Net Content: **ONE Gallon (3.78L)**

K20Lot: -2020

Expiration: -2022