

See package insert for complete product information.

DOSAGE AND ADMINISTRATION: IMPROVEST should be administered via subcutaneous injection into the post auricular region of the neck. Each intact male pig should receive two 2-mL doses. The first dose should be administered no earlier than 9 weeks of age. The second dose should be administered at least 4 weeks after the first dose. Pigs should be slaughtered no earlier than 4 weeks and no later than 8 weeks after the second dose. In case of misdosing, the animal should be re-dosed immediately.

WITHDRAWAL PERIODS: No withdrawal period is required when used according to labeling.

WARNINGS: Not for human use. Keep out of reach of children. Pregnant women should not administer this product. Women of childbearing age should exercise extreme caution when handling this product. In case of accidental self-injection seek immediate medical attention, as this product could negatively affect reproductive physiology in men and women. People previously injected with IMPROVEST should not administer the product because of a booster effect of the second dose. Read package insert carefully prior to use.

Store under refrigeration at 2°-8°C (36°-46°F). Once broached, product may be stored under refrigeration for 28 days. Store bottles in carton until used. Protect from light. Protect from freezing.

Improv^{est}[®]

(Gonadotropin Releasing Factor Analog-Diphtheria Toxoid Conjugate, 0.2 mg/mL)

Sterile Solution

For subcutaneous injection in intact male pigs intended for slaughter.

For use in animals only

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

mL (doses, mL each)

NADA #141-322, Approved by FDA



Dist. by: Pfizer Animal Health
Div. of Pfizer Inc., New York,
NY 10017 1534000

Lot/Exp:

IMPRINT AREA:
NO VARNISH, NO COPY, NO INK