

For Barn and Farm Use

This product cleans by removing dirt, grime, mildew, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine or poultry facilities, and farms. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

This product cleans, disinfects and deodorizes in one step. Its non-abrasive formula is designed for use on hard, non-porous surfaces: plated or stainless steel, aluminum, chrome, glazed porcelain, glazed tile, sealed concrete, laminated surfaces associated with floors, walls, non-food contact countertops, animal equipment found in barns, pens and stalls and animal housing facilities.

This product is designed for use as a cleaner, disinfectant, sanitizer on hard non-porous surfaces of vehicles, trucks, trailers, carts, animal crates, grooming equipment, handling equipment, buckets, brushes, tables, transportation crates, foot baths, automatic feeders, troughs, feed racks, wash stalls, non porous fencing, gates, gathering guiding / controlling equipment, and non porous harnesses and leashes; commonly found in animal housing facilities, poultry, stall, barn, pen, agricultural and livestock buildings, and breeding facilities.

To Refill Concentrate From Large Containers Into Smaller Containers: This product may be used to fill and refill clean, properly labeled containers for dilution elsewhere within your facility. Make sure the small container has been cleaned, dried and properly labeled. Also make sure other items (funnels or hand pumps) are properly cleaned and dried. To refill, simply pour or pump product from the larger container directly into the smaller one being careful not to spill any product. Keep both containers sealed when not in use.

Modes of Application

This product can be applied by mop, sponge, cloth, paper towel, coarse trigger sprayer, or auto-scrubber. Change cloth, sponges or towels frequently to avoid redeposition of soil.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not recommended for use on copper, brass or marble.

To Prepare Use Solution:

Ratio	Per Gallon of Water	Per Liter of Water
1:16	8 oz.	64 mL
1:32	4 oz.	32 mL
1:64	2 oz.	16 mL
1:128	1 oz.	8 mL
1:256	0.5 oz.	4 mL

For Use as a Daily One-Step Cleaner / Disinfectant: Dilute at 8.0 oz. of product per gallon of water (1:16).

- Pre-clean heavily soiled areas.
- Apply Use Solution by hand pump coarse trigger sprayer to hard, non-porous environmental surfaces. Spray 6-8 inches from surface; making sure to wet surfaces thoroughly.
- All surfaces must remain wet for 5 minutes.
- Wipe surfaces and let air dry.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16).

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry.

For Use as a Cleaner / Disinfectant in non-food handling areas of Food Processing Plants: Dilute at 8.0 oz. of product per gallon of water (1:16).

- Apply Use Solution evenly over surface. Be sure to wet all surfaces thoroughly.
- Allow product to remain on surface for 5 minutes.
- Wipe with clean cloth, sponge or paper towel.
- For heavily soiled areas, thoroughly clean surface prior to disinfecting.

For Treatment of Animal Housing Facilities including Poultry barns: Dilute at 8.0 oz. of product per gallon of water (1:16).

- Remove all animals and feeds from areas being treated.
- Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
- Empty or cover all troughs, racks and other feeding and watering appliances.
- Thoroughly clean all surfaces with soap or detergent and rinse with water.
- Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for 5 minutes.
- Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
- Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried.
- For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand 5 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

For use as a Boot / Shoe Wash: Dilute at 8.0 oz. of product per gallon of water (1:16).

To prevent cross contamination into animal areas, shoe baths containing one inch of freshly made disinfecting solution must be placed at all entrances to building. Scrape waterproof shoes or boots and thoroughly scrub with this product and rinse with water. Place in INTERvention Farm Animal Care Disinfectant Cleaner & Deodorizer. Allow product to remain in contact with shoes or boots for 5 minutes and allow to air dry. All treated surfaces that will contact feed or drinking water must be thoroughly scrubbed with soap or detergent and then rinsed with potable water before reuse.

INTERvention®

EPA Reg. No.
74559-4

EPA Est. No.
74559-CAN-1

Part No. 24309

INTERvention®

Farm Animal Care

Disinfectant Cleaner & Deodorizer Concentrate

ACTIVE INGREDIENT
Hydrogen Peroxide 4.25%
OTHER INGREDIENTS 95.75%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Additional Precautionary Statements on Side Panel of label.

See reference sheet for a complete list of pathogenic organisms, additional features, claims, and directions for use.

FOR BARN AND FARM USE

Fragrance Free Formula



Net Contents: 5 Gallon (18.9L) **5G**

For Barn and Farm Use

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

SMALL NONREFILLABLE CONTAINERS: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¾ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Precautionary Statements

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

Claims

See reference sheet for a complete list of organisms.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Bacteria:

Pseudomonas aeruginosa (ATCC 15442)
Salmonella enterica (ATCC 10708) (formerly known as *Salmonella choleraesuis*)
Staphylococcus aureus (ATCC 6538)
Escherichia coli 0157:H7 (ATCC 35150)

Listeria monocytogenes (ATCC 19111)
Streptococcus pyogenes (ATCC 12204)

Antibiotic-Resistant Bacteria:

Staphylococcus epidermidis (MRSE) (ATCC 51625)
Staphylococcus aureus (MRSA) (ATCC 33591)
Streptococcus pneumoniae, Penicillin Resistant (PRSP) (ATCC 700671)
Enterococcus faecium, Resistant to Vancomycin (VRE) (ATCC 51559)
Klebsiella pneumoniae, Carbapenem-resistant (KPC), (ATCC BAA-1705)

Fungi:

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Trichophyton mentagrophytes (ATCC 9533)

Microsporum canis (ATCC 10214)

*Viruses:

At 1:16 - 1:64 dilution (2 - 8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

*Influenza Type A2 (Hong Kong) (VR-544)

*Norovirus (Feline Calicivirus as the surrogate) (VR-782)

*Human Coronavirus (VR-740)

*Parainfluenza Type 3 (VR-93)

*Rhinovirus Type 37, Strain 151-1 (ATCC VR-1147)

*Vaccinia Virus (smallpox vaccine virus) (ATCC VR-119)

*Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swine flu).

Veterinary *Viruses:

*Duck Hepatitis B Virus (DHBV)

*Bovine Viral Diarrhea Virus (BVDV) (ATCC VR-1422)

*Avian Influenza A (H3N2) (ATCC VR-2072)

*Equine Influenza A (H3N8) virus (strain Alaska/A/EQ/AK/29759/91)

*Porcine Respiratory & Reproductive Syndrome (PRRS) virus (Strain NVSL)

*Equine Herpes virus type 1 (strain EH1)

*Porcine Epidemic Diarrhea virus (Strain Colorado 2013 Isolate)

Bactericidal Stability of Use-Solution – This product, when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed. If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Escherichia coli (ATCC 11229)

Listeria monocytogenes (ATCC 19111)

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708) (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Malodor Activity – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors, when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water in 30 seconds or when used as directed at a 1:128 dilution (1 oz. per gallon of water), in the presence of 200 ppm hard water in 3 minutes.

*Virucidal: 5 Min. @ 1:64

Bactericidal: 5 Min. @ 1:16

Fungicidal: 5 Min. @ 1:16

Broad-Spectrum
Non-Food Contact Sanitizer:
3 Min. @ 1:128

SPOT VARNISH

READ S.D.S. BEFORE USING PRODUCT

SDS available online at www.virox.com

Si no puede leer en ingles, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto.



Manufactured by:

Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6M1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50105-5

Made in Canada

VAH023 (15/349)