



COMMITTED TO QUALITY. DEDICATED TO ANIMAL HEALTH.

Ivermax[®]

(ivermectin topical solution) Pour-on for Cattle



PRODUCT DESCRIPTION:

This pour-on formulation of ivermectin provides effective treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle. Contains 5mg of ivermectin per milliliter.

SIZES:

- 250 mL (8.5 fl oz)
- 1 L (33.8 fl oz)
- 2.5 L (84.5 fl oz)
- 5 L (169 fl oz)
- 20 L (676 fl oz)

Store away from excessive heat (104°F/40°C) and protect from light.

See package insert or label for complete indications, precautions, warnings, residue information, and use directions.



For More Information:
info@aspenveterinaryresources.com

Ivermax[®]

(ivermectin topical solution) Pour-on for Cattle

PACKAGING LABEL:



NDC # 46066-751-03

aspen
VETERINARY RESOURCES, LTD.

IVERMAX[®]

(ivermectin topical solution)
POUR-ON FOR CATTLE

Parasiticide • Contains 5 mg ivermectin/mL

Kills: Roundworms (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies

Approved by FDA under ANADA # 200-318

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS: For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

See package insert for the complete indications and use directions.

DOSEAGE: The dose rate is 1 mL per 22 lb of body weight. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead. Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

ADMINISTRATION: Use dosing equipment (applicator, draw-off tubing, cap with stem) compatible with Ivermax[®] (ivermectin topical solution) Pour-On. Other dosing equipment may be incompatible, resulting in leaking, incorrect dosage and leakage. Follow the dosing equipment manufacturer's directions for adjusting the dose and proper use and maintenance of the dosing equipment. Connect the dosing applicator and the draw-off tubing to the container as follows:

- Attach the open end of the draw-off tubing to an appropriate dosing applicator. Attach draw-off tubing to the cap with the stem. Replace the shipping cap with the cap having the draw-off tubing. Gently prime the dosing applicator, checking for leaks.
- When the interval between uses of the applicator gun is expected to exceed 12 hours, disconnect the gun, draw-off tubing and cap with stem from the product container, empty the product from the gun and bubble back into the product container and replace with the shipping cap.

ANIMAL SAFETY: Studies conducted in the U.S.A. have demonstrated the safety margin for ivermectin. Based on plasma levels, the typically applied formulation is expected to be at least as well tolerated by breeding animals as is the substantially formulated which had no effect on breeding performance.

WARNING:

NOT FOR USE BY HUMANS. Keep this and all drugs out of the reach of children. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-724-6375 or online at www.fda.gov/oc/medwatch.

WARNING: FLAMMABLE! KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME AND OTHER SOURCES OF IGNITION.

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

REQUIRE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-weaning calves. Do not use in calves to be processed for veal.

PRECAUTIONS: Store at 20°C - 25°C (68°F - 77°F) and protect from light. Excursions permitted to 15°C (59°F).

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced. Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally. Opalescence in the formulation may occur when Ivermax[®] (ivermectin topical solution) Pour-On is stored at temperatures below 32°F. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g., caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermax[®] (ivermectin topical solution) Pour-On is not recommended for use in species other than cattle. Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Restricted Drug (California) - Use only as directed

Manufactured For:
Aspen Veterinary Resources,® Ltd.
Liberty, MO 64068, USA
www.aspenveterinaryresources.com

A187B
I#V6332 #I#V6338 05/20

LOT:
EXP:

0 79335 01090 1

Net Contents:
20 Liter (676 fl oz)
Contains 800-550 lb Doses

Product Code	Product Size	Case
15703301	250 mL	12/Case
15893311	1 L	12/Case
15998644	2.5 L	4/Case
14563321	2.5 L	4/Case
21271833	2.5 L	By the Case*
21263876	2.5 L	By the Case*
16998644	5 L	2/Case
21270267	5 L	By the Case*
16998644	20 L	1/Case