

INDICATIONS: For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.

DOSAGE AND ADMINISTRATION: May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique.

Cattle and Horses: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

Swine and Sheep: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

If administered subcutaneously divide the dosage into several sites of injection and massage points of injection to aid in absorption and help prevent inflammation and/or sloughing.

Store between 15°C and 30°C (59°F and 86°F)

Rev. 04-16



® Registered trademark of Agri Laboratories, Ltd.

**Manufactured for
Agri Laboratories, Ltd.
St. Joseph, MO 64503**

LOT

EXP



NDC 57561-803-60

Lactated Ringers Injection

STERILE NONPYROGENIC SOLUTION

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For Animal Use Only

**Keep Out of Reach
of Children**

NET CONTENTS: 1000 mL



 **AgriLabs®**
Making A Healthy Difference®

Each 100 mL contains:
Sodium Chloride 600 mg
Sodium Lactate 310 mg
Potassium Chloride 30 mg
Calcium Chloride Dihydrate 20 mg
Water for Injection q.s.

The Calcium, Potassium and Sodium contents are approximately 2.7, 4.0, and 130 mEq/liter, respectively. Total Osmolar Concentration: 269 mOsm per liter (calculated).

CAUTION: Solution should be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. Use entire contents when first opened.

WARNING: Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis.

18-803-60

RMS 92-579

