

For Intramuscular Use in Swine.
Restricted Drug (California) - Use Only as Directed.
For Use in Animals Only.

DOSAGE: Usual daily dose for arthritis or mycoplasma pneumonia - 5 mg per pound of body weight (1 mL per each 60 pounds of body weight) intramuscularly for three to seven days. See package insert for complete product information.

Residue Warning: Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

Human Warnings: Not for human use. Keep out of reach of children.

Store at controlled room temperature between 20°-25°C (68° to 77°F) [see USP]. KEEP FROM FREEZING.

Product No. 1LIN020 8LIN002A

LincoMed[®] 300

lincomycin injection, USP



Swine Antibiotic

ANADA 200-368, Approved by FDA

Net: 100 mL (3.3 fl oz)

Manufactured for:
Bimeda, Inc.
Le Sueur, MN 56058



Bimeda

Contains per mL: Lincomycin hydrochloride equivalent to lincomycin 300 mg; also benzyl alcohol 9.45 mg added as a preservative.

INDICATIONS: Indicated for the treatment of arthritis caused by susceptible organisms and for mycoplasma pneumonia.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Manufactured by: Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda, Inc. and Bimeda-MTC Animal Health Inc.
are Divisions of Cross Vetpharm Group Ltd.

8 99855 60002 8

LincoMed is a Registered Trademark of Bimeda, Inc.