

# LINCOMIX® 20 (lincomycin)

## Type A Medicated Article 20 grams/lb

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*.

For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*.

### Each pound contains:

#### Active Drug Ingredient

Lincomycin ..... 20 grams  
(as lincomycin hydrochloride agricultural grade)

#### Inactive Ingredients

Soybean hulls, #20 grind; light liquid paraffin, Ph.Eur.

#### IMPORTANT:

**Must be thoroughly mixed in feeds before use.  
Store opened bag in dry place to prevent caking.  
Store at room temperature.**

Approved by FDA under NADA # 097-505

LOT:  
EXP:  
MFG:

## NET WEIGHT 50 lb (22.7 kg)

LINCOMIX® 20  
(lincomycin)  
Type A Medicated Article  
20 grams/lb

TAKE TIME



OBSERVE  
LABEL  
DIRECTIONS

LINCOMIX® 20  
(lincomycin)  
Type A Medicated Article  
20 grams/lb

# LINCOMIX® 20 (lincomycin)

## Type A Medicated Article 20 grams/lb



#### BROILERS

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

#### DIRECTIONS FOR USE

**For the control of necrotic enteritis:** LINCOMIX 20, 20 grams/lb, should be mixed into the complete feed supplied to broiler chickens so that the final feed contains 2 grams of lincomycin per ton of feed.

#### MIXING DIRECTIONS

##### Type C Medicated Feeds

Intermediate Premix Amount of LINCOMIX 20 per 1000 lb (454 kg) of Feed Ingredients	Complete Feed Amount of Intermediate Premix to Provide 2 Grams of lincomycin per ton of Type C Medicated Feed
	<b>lincomycin per ton of feed</b>
	<b>2 grams</b>
50 lbs	2 lbs
10 lbs	10 lbs
5 lbs	20 lbs

**WARNING:** When using LINCOMIX 20 in approved combinations with other drugs, follow the required withdrawal times for those drugs. No drug withdrawal period is required before slaughter of birds fed LINCOMIX 20 at the approved concentration (2 grams lincomycin per ton of feed).

**CAUTION:** Not for use in layers, breeders, or turkeys. (For additional **Cautions**, see below.)

#### SWINE

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*. For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*.

#### DIRECTIONS FOR USE

**For the treatment of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks, or until signs of disease (watery, mucoid or bloody stools) disappear.

**For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks, or until signs of disease (watery, mucoid or bloody stools) disappear, followed by 40 grams of lincomycin per ton.

**For the control of swine dysentery and porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** Feed 40 grams of lincomycin per ton of complete feed as the sole ration. For use in animals or on premises with a history of swine dysentery, but where symptoms have not yet occurred.

**For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*:** Feed 100-200 grams of lincomycin per ton of complete feed as the sole ration for 21 days.

#### MIXING DIRECTIONS

##### Type C Medicated Feeds

**For treatment of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** To make complete feed containing 100 grams of lincomycin, add 5 lbs of LINCOMIX 20 per ton.

**For control of swine dysentery and porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** To make complete feed containing 40 grams of lincomycin, add 2 lbs of LINCOMIX 20 per ton.

**For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*:** To make complete feed containing 100-200 grams of lincomycin, add 5-10 lbs of LINCOMIX 20 per ton.

Additional mixing directions to make complete feed containing 40, 100 or 200 grams of lincomycin per ton are provided below.

#### MIXING DIRECTIONS

Intermediate Premix Amount of LINCOMIX 20 per 1000 lb (454 kg) of Feed Ingredients	Complete Feed Grams of lincomycin per ton of Type C Medicated Feed
	<b>lincomycin per ton of feed</b>
	<b>40 grams</b> <b>100 grams</b> <b>200 grams</b>
125 lbs	16 lbs    40 lbs    80 lbs
100 lbs	20 lbs    50 lbs    100 lbs
50 lbs	40 lbs    100 lbs    200 lbs

**WARNING:** When using LINCOMIX 20 in approved combinations with other drugs, follow the required withdrawal times for those drugs. No drug withdrawal period is required before slaughter of swine fed LINCOMIX 20 at approved concentrations (40, 100 or 200 grams lincomycin per ton of feed).

#### NOT FOR HUMAN USE

**CAUTION:** Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the snout. On rare occasions, some pigs may show reddening of the skin and irritate behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Good Manufacturing Practices Should Be Observed in Preparing Feeds Containing LINCOMIX 20. This Includes Appropriate Clean-out Procedures to Avoid Cross-Contamination.

Restricted Drug (California)—Use Only as Directed

Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007

Made in China

501917  
40027872