

NO COPY

SWINE: Directions for use
DOSAGE AND ADMINISTRATION
Dosage: Administer at a dose rate of 250 mg of lincomycin per gallon of drinking water. In clinical studies, this dose rate provided an average of 3.8 mg of lincomycin per lb of body weight per day.
Treatment Period: The drug should be administered for a minimum of 5 consecutive days beyond the disappearance of symptoms (bloody stools) up to a maximum of 10 consecutive days. If water treatment is discontinued prior to this time, a lincomycin treatment program may be continued with lincomycin premix at 100 g lincomycin per ton of complete feed as the sole ration according to label directions.
Administration: This packet will medicate 128 gallons of drinking water providing 250 mg/gallon. A dose of 3.8 mg lincomycin per lb of body weight may be maintained by medicating the drinking water at a concentration of 250 mg per gallon of drinking water when pigs are consuming 1.5 gallons per 100 lbs of body weight per day. Under these circumstances the concentration of lincomycin required in medicated water may be adjusted to compensate for variations in age and weight of animals, the nature and severity of disease symptoms, environmental temperature and humidity, each of which affects water consumption. For use in automatic water proportioners, prepare the stock solution by dissolving 1 packet in 1 gallon of water; then adjust the proportioner to deliver 1 oz of stock solution per gallon of drinking water.

CORNER RADIUS

Lincomycin Hydrochloride Soluble Powder

Antibacterial

This packet contains as an active ingredient:
 Lincomycin HCl, equivalent to 32 g lincomycin
 Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).
 See back for instructions.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Net Weight
80 grams
(2.82 oz)

For Oral Use in Swine and Broiler Chickens Only
Keep Out of Reach of Children

INDICATIONS AND USAGE:

Swine: Lincomycin Hydrochloride Soluble Powder is indicated for the treatment of swine dysentery (bloody scours).
Broiler Chickens: Lincomycin Hydrochloride Soluble Powder is indicated for the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.

RESIDUE WARNINGS:

No drug withdrawal period is required before slaughter of swine receiving Lincomycin Hydrochloride Soluble Powder at the approved level of 250 mg per gallon of drinking water, nor before slaughter of birds receiving Lincomycin Hydrochloride Soluble Powder at the approved level of 64 mg per gallon of drinking water.



Distributed by:
 Huvepharma, Inc.
 Peachtree City, GA 30269

Restricted Drug (California) - Use only as directed.
 Not for Human Use.
 ANADA 200-189, Approved by FDA
 13899200

HUVEPHARMA is a registered trademark of Huvepharma EOOD.

CORNER RADIUS

BROILER CHICKENS: Directions for use
DOSAGE AND ADMINISTRATION
Dosage: Administer at a dose rate of 64 mg of lincomycin per gallon of drinking water.
Treatment Period: Start medication as soon as the diagnosis of necrotic enteritis is determined. If improvement is not noted within 24 to 48 hours, consult a licensed veterinarian or veterinary diagnostic laboratory to determine diagnosis. The drug should be administered for 7 consecutive days.
Administration: This packet will medicate 500 gallons of drinking water providing 64 mg/gallon.
PRECAUTIONS:
 Discard medicated drinking water if not used within 2 days. Fresh stock solution should be prepared daily. Do not use the water treatment and the feed treatment simultaneously. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to water containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.
Swine: If clinical signs of bloody scours (watery, mucoid or bloody stools) have not improved during the first 6 days of medication, discontinue treatment and redetermine the diagnosis. Occasionally, swine fed lincomycin may within the first 2 days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5 to 8 days without discontinuing the lincomycin treatment. The safety of lincomycin has not been demonstrated for pregnant swine or swine intended for breeding.
Chickens: Not for use in layer and breeder chickens.

NO COPY

Time/Date: 13899200.pdf 1 7/7/15 1:36 PM

Project No.	Artwork Number	Description	Country
11422	13899200	Lincomycin Soluble	US
Dimensions		Drawing No.	SKU No.
5.498" x 11.625"		DWG-102294-01	3765000
Additional Info:			Item
JDE Component Code - 3902BF Substrate: Aluminum			Foil
Colors: Black PMS 307 PMS 1485 Printable area			<input type="checkbox"/> EDITOR'S COPY GS: DATE:
Mgr	Rev	GA	PR
GS	6		
GA			
		CHANGES OK	CHANGES OK
		GS / ART REV (LCA)	GS / ART REV (FA)
		CHANGES OK	CHANGES OK