

4.5"



LinxMed-SP[®] (lincomycin hydrochloride) Soluble Powder

FOR THE TREATMENT OF DYSENTERY IN SWINE AND FOR THE CONTROL OF NECROTIC ENTERITIS IN BROILER CHICKENS

Antibacterial

CAUTION: Federal (US) law restricts this drug to use by or on the order of a licensed veterinarian.

For oral use in swine and broiler chickens only. Not for use in layer and breeder chickens.

This packet contains as active ingredient:

Lincomycin hydrochloride

Equivalent to lincomycin 16.0 g

FOR ANIMAL USE ONLY

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

ANADA 200-377, Approved by FDA



Net Weight: 40 g (1.41 oz)

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

**2" gap
bottom of pouch**

Product No. TLIN004
8LIN005 Rev. 08/16



Oakbrook Terrace, IL 60181
LinxMed-SP[®] is a Registered Trademark of Bimeda, Inc.
N. A. Corp. Address: Bimeda, Inc., One Tower Lane.

Store at controlled room temperature 20°C to 25°C (68°F to 77°F).

No drug withdrawal period is required before slaughter of swine receiving LinxMed-SP[®] Soluble Powder at the approved level of 250 mg per gallon of drinking water, nor before slaughter of birds receiving the approved level of 64 mg per gallon of drinking water.

RESIDUE WARNINGS:

1. Discard medicated drinking water if not used within 2 days. Fresh stock solution should be prepared daily.
2. If clinical signs of bloody scours (watery, mucoid or bloody stools) have not improved during the first 6 days of medication, discontinue treatment and re-determine the diagnosis.
3. Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have not been demonstrated for pregnant swine or swine intended for breeding.
4. The safety of lincomycin has not been demonstrated for pregnant swine or swine intended for breeding.
5. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to water containing lincomycin treatment. Ingestion by these species may result in severe gastrointestinal effects.
6. Do not use the water treatment and the feed treatments simultaneously.
7. Not for use in layer and breeder chickens.

PRECAUTIONS

feeding lincomycin premix at 2 grams lincomycin per ton of complete feed.

NOTE: After water medication is discontinued, a recommended control program for necrotic enteritis consists of 250 mg per gallon of drinking water providing 64 mg/gallon.

ADMINISTRATION: This packet will medicate 250 gallons of drinking water providing 64 mg/gallon. Should be administered for 7 consecutive days.

TREATMENT PERIOD: Start medication as soon as the diagnosis of necrotic enteritis is determined. The drug dosage: Administer at a dose rate of 64 mg of lincomycin per gallon of drinking water.

INDICATIONS: LinxMed-SP[®] Soluble Powder is indicated for the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.

BROILER CHICKENS: DIRECTIONS FOR USE

feeding lincomycin premix at 40 grams lincomycin per ton of complete feed as the sole ration.

NOTE: After a treatment program is discontinued, a control program for swine dysentery may be followed by 250 mg per gallon of drinking water.

proportions, prepare the stock solution by dissolving two packets in one gallon of water, then adjust the environmental temperature and humidity, each of which affects water consumption. For use in automatic water dispensers to compensate for variations in age and weight of animals, the nature and severity of disease symptoms, weight per day. Under these circumstances the concentration of lincomycin required in medicated water may be adjusted to compensate for variations in age and weight of animals, the nature and severity of disease symptoms, weight per day.

ADMINISTRATION: This packet will medicate 64 gallons of drinking water providing 250 mg/gallon. A dose of 3.8 mg lincomycin per pound of body weight may be maintained by medicating 250 mg/gallon. A dose of 3.8 mg lincomycin per pound of body weight may be maintained by medicating 250 mg/gallon. A dose of 3.8 mg lincomycin per pound of body weight may be maintained by medicating 250 mg/gallon. A dose of 3.8 mg lincomycin per pound of body weight may be maintained by medicating 250 mg/gallon.

TREATMENT PERIOD: The drug should be administered for a minimum of 5 consecutive days beyond disappearance of symptoms (bloody stools) up to a maximum of 10 consecutive days. If water treatment is discontinued prior to this time, a lincomycin treatment program may be continued with lincomycin premix at 100 mg per gallon of drinking water.

INDICATIONS: LinxMed-SP[®] Soluble Powder is indicated for the treatment of swine dysentery (bloody scours), dysentery, and necrotic enteritis in broiler chickens.

FOR THE TREATMENT OF DYSENTERY IN SWINE AND FOR THE CONTROL OF NECROTIC ENTERITIS IN BROILER CHICKENS

LinxMed-SP[®]
(Lincomycin hydrochloride)
Soluble Powder