



LinxMed-SP[®] (lincomycin hydrochloride) Soluble Powder

FOR THE TREATMENT OF DYSENTERY IN SWINE AND FOR THE CONTROL OF NECROTIC ENTERITIS IN BROILER CHICKENS

Antibacterial

CAUTION: Federal (US) law restricts this drug to use by or on the order of a licensed veterinarian.

**For oral use in swine and broiler chickens only.
Not for use in layer and breeder chickens.**

This packet contains as active ingredient:

Lincomycin hydrochloride

Equivalent to lincomycin 192 g

**FOR ANIMAL USE ONLY
NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

Restricted Drug (California) - Use Only as Directed
ANADA 200-377, Approved by FDA



**Net Weight:
480 g (1 lb. 0.93 oz)**

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com



Product No: 1LIN014
8LIN020 Rev. 08/16



TAKE TIME



Store at controlled room temperature 20°C to 25°C (68°F to 77°F).
LinxMed-SP[®] is a Registered Trademark of Bimeda, Inc.
N.A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181

RESIDUE WARNINGS:
No drug withdrawal period is required before slaughter of birds receiving LinxMed-SP[®] Soluble Powder at the approved level of 64 mg per gallon of drinking water.
250 mg per gallon of drinking water, nor before slaughter of swine receiving LinxMed-SP[®] Soluble Powder at the approved level of 64 mg per gallon of drinking water.

1. Discard medicated drinking water if not used within 2 days. Fresh stock solution should be prepared daily.
2. If clinical signs of bloody scours (watery, mucoid or bloody stools) have not improved during the first 5 days of medication, discontinue treatment and redetermine the diagnosis.
3. Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment.
4. The safety of lincomycin has not been demonstrated for pregnant swine or swine intended for breeding.
5. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to water containing lincomycin treatment. Ingestion by these species may result in severe gastrointestinal effects.
6. Do not use the water treatment and the feed treatments simultaneously.
7. Not for use in layer and breeder chickens.

PRECAUTIONS

INDICATIONS: LinxMed-SP[®] Soluble Powder is indicated for the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.

DOSEAGE: Administer at a dose rate of 64 mg of lincomycin per gallon of drinking water.
TREATMENT PERIOD: Start medication as soon as the diagnosis of necrotic enteritis is determined. The drug should be administered for 7 consecutive days.

ADMINISTRATION: This packet will medicate 3,000 gallons of drinking water providing 64 mg/gallon.
NOTE: After a treatment program is discontinued, a control program for necrotic enteritis consists of feeding lincomycin premix at 2 grams lincomycin per ton of complete feed.

BROILER CHICKENS: DIRECTIONS FOR USE

40 grams lincomycin per ton of complete feed as the sole ration.
NOTE: After a treatment program is discontinued, a control program for swine dysentery may be followed by feeding lincomycin premix at 40 grams lincomycin per ton of complete feed.

ADMINISTRATION: This packet will medicate 768 gallons of drinking water providing 250 mg/gallon. A dose of 3.8 mg lincomycin per pound of body weight may be maintained by medicating the drinking water at a concentration of 250 mg per gallon of drinking water per pig are consuming 1.5 gallons per 100 lbs of body weight per day. Under these circumstances the concentration of lincomycin required in medicated water may be adjusted to compensate for variations in age and weight of animals, the nature and severity of disease symptoms, environmental temperature and humidity, each of which affects water consumption. For use in automatic water proportioners, prepare the stock solution by dissolving one packet in six gallons of water, then adjust the proportioner to deliver 1 ounce of stock solution per gallon of drinking water.

TREATMENT PERIOD: The drug should be administered for a minimum of 5 consecutive days beyond disappearance of symptoms (bloody stools) up to a maximum of 10 consecutive days. If water treatment is discontinued prior to this time, a lincomycin treatment program may be continued with lincomycin premix at 100 grams lincomycin per ton of complete feed as the sole ration according to label directions.

DOSEAGE: Administer at a dose rate of 250 mg of lincomycin per gallon of drinking water. In clinical studies, this dose rate provided an average of 3.8 mg of lincomycin per pound of body weight per day.

SWINE: DIRECTIONS FOR USE

INDICATIONS: LinxMed-SP[®] Soluble Powder is indicated for the treatment of swine dysentery (bloody scours).
DOSEAGE: Administer at a dose rate of 250 mg of lincomycin per gallon of drinking water. In clinical studies, this dose rate provided an average of 3.8 mg of lincomycin per pound of body weight per day.

FOR THE TREATMENT OF DYSENTERY IN SWINE AND FOR THE CONTROL OF NECROTIC ENTERITIS IN BROILER CHICKENS

LinxMed-SP[®] Soluble Powder
(lincomycin hydrochloride)

19.0000"

7.0000"