

4697



Liquamycin[®] LA-200[®] (oxytetracycline injection)

Antibiotic

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.

For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine

For animal use only

Net Contents: 500 mL

NADA #113-232, Approved by FDA



4697



Liquamycin[®] LA-200[®] (oxytetracycline injection)

Antibiotic

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.

For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine

For animal use only

Net Contents: 500 mL

NADA #113-232, Approved by FDA

DESCRIPTION
Liquamycin LA-200 is a sterile, preservative-free solution of the antibiotic oxytetracycline hydrochloride. Each mL contains 200 mg of oxytetracycline hydrochloride (equivalent to 180 mg of oxytetracycline base), 4.2% benzalkonium chloride, 5.0% povidone-iodine, and 0.1% sodium metabisulfite (as a preservative). The solution is adjusted to a pH of 7.0 to 8.0.

INDICATIONS AND USAGE
Liquamycin LA-200 is indicated for the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.

Directions
Injectable: 200 mg (0.5 mL) per lb of body weight per day for 14 days. For calves, injectable may be administered intramuscularly or subcutaneously. For adult cattle, injectable may be administered intramuscularly. For swine, injectable may be administered intramuscularly. For preruminating calves, injectable may be administered intramuscularly. For adult cattle, injectable may be administered intramuscularly. For swine, injectable may be administered intramuscularly. For preruminating calves, injectable may be administered intramuscularly.

Warnings
Liquamycin LA-200 is contraindicated in animals with known hypersensitivity to tetracyclines. It is also contraindicated in animals with known hypersensitivity to any of the components of this product. It is contraindicated in animals with known hypersensitivity to any of the components of this product.

Adverse Reactions
The most common adverse reactions reported in animals treated with Liquamycin LA-200 are diarrhea, anorexia, and depression. Other adverse reactions include skin reactions, photosensitivity, and bone marrow depression.

Drug Interactions
Liquamycin LA-200 may interact with other drugs, including antacids, iron supplements, and certain antibiotics. Consult your veterinarian for more information.

How Supplied
Liquamycin LA-200 is available in 500 mL glass vials.

Storage and Stability
Store at 20° to 25° (68° to 77°) F. Excursions permitted to 15° to 30° (59° to 86°) F. Keep from light.

Animal Health
NADA #113-232
79-4864-00-1
Made in USA

FOR ANIMAL USE ONLY