

AmerisourceBergen

MWI Animal Health®

Safe-Guard® Swine Dewormer 1.80%

FENBENDAZOLE TYPE B MEDICATED FEED

For the further manufacture of swine feed

INDICATIONS

For the treatment and control of: **Lungworms:** Adult *Metastrongylus apri*, Adult *Metastrongylus. pudendotectus*; **Gastrointestinal worms:** Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworms (*Ascaris suum*), Adult nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*), Adult small stomach worms (*Hyoststrongylus rubidus*), Adult and larvae (L2, L3, L4 stages--intestinal mucosal forms) whipworms (*Trichuris suis*), and **Kidney Worms:** Adult and larvae *Stephanurus dentatus*.

ACTIVE DRUG INGREDIENT

Fenbendazole1.80% (8.164 grams/lb)

INGREDIENTS

Calcium Carbonate, Roughage Products, Molasses Products, Soybean Oil.

MIXING AND USE DIRECTIONS

Mix 36.75 pounds of the 1.8% Type B Medicated Feed with 1963.25 pounds of feed ingredients to manufacture Type C medicated feed containing 300 of fenbendazole per ton.

The resulting Type C medicated feed is to be fed as the sole ration for 3 to 12 consecutive days at the rate of 0.91 to 6.80 pounds daily per 100 pounds of body weight to provide a total dose of 4.08 milligrams fenbendazole per pound of body weight.

Type C Medicated Feed – Feeding Directions (300 g/t FBZ).

Pig Wt. (lbs)	Days Type C medicated feed are fed (lbs/d)		
	3	6	12
50	0.5	0.23	0.11
100	0.9	0.45	0.23
300	2.7	1.36	0.68

Feeding a 300 lb gilt for 3 consecutive days provides 0.8 ounces of the 1.8% Type B Medicated Feed. A 400 lb sow would consume 1.2 ounces of the Type B Medicated Feed in a 3 consecutive day feeding period.

WITHDRAWAL PERIODS

Swine must not be slaughtered for human consumption
Within 4 days following last treatment with this drug product.

OTHER WARNINGS

Parasite resistance may develop to any wormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

NOTE: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Manufactured for
MWI Animal Health
Boise, ID 83705

Lot



21784457255

Safe-Guard® is a registered trademark of Intervet International B.V.

NET WEIGHT: 20 LBS (9.07 kg)

Approved by FDA under NADA # 131-675

11034173

Lt Blue

Rev 071822