



NeoMed[®] 325 (neomycin sulfate) Commercial Grade Soluble Powder

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

EACH PACKET CONTAINS: 142.8 g neomycin sulfate (commercial grade) equivalent to 100 g neomycin.

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

Restricted Drug (California) - Use Only as Directed
ANADA 200-050, Approved by FDA



Net Weight: 200 g (7 oz)



Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

4" gap bottom of pouch



NeoMed[®] is a registered trademark of Bimeda, Inc.
N.A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181
Product No. 1NEC010
8NEC008 Rev. 06/16



When storing partially used packet, securely close to protect from moisture. Store in a dry place.
Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

RESIDUE WARNINGS: Discontinue treatment prior to slaughter by at least the number of days listed below for appropriate species:
Cattle: 1 day
Sheep: 2 days
Swine and goats: 3 days
Turkeys: 0 days
A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

IMPORTANT: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

PRECAUTIONS: To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.
If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

DRINKING WATER: Swine - Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one gallon.
Level Teaspoonful = US Standard Measure

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoonful in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Head/Flock Treatment: Each packet will treat 14,300 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 14,300 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

DOSE AND ADMINISTRATION: Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Add to drinking water - Not for use in liquid supplements.

Fig's Weighing 25 to 50 Pounds: 1 Packet
Fig's Weighing 50 to 100 Pounds: 1.5 Packets
Fig's Weighing Over 100 Pounds: 2 Packets

NeoMed[®] 325 (neomycin sulfate) Commercial Grade Soluble Powder