



# 11 pounds

Net Weight (5 kg)

 **NOVARTIS**  
ANIMAL HEALTH



**NEPOREX**<sup>®</sup>  
**2 SG**

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN INJURY, POOR FLY CONTROL, AND/OR ILLEGAL RESIDUES.**

Do not apply this product in such a manner as to directly or through spray drift expose workers or other persons.

**For Housefly, Lesser Housefly, Stable Fly and Soldier Fly Control in Cattle Operations, Hog Operations, Poultry Operations Including Layer and Breeder Chickens, and Horse Barns Where Horses are not Intended for Slaughter or Food**

Fly control in cattle, hog, poultry, and horse operations should include appropriate sanitary and management practices to reduce the number and size of fly breeding sites. A successful sanitary and management program may allow less than constant use of insecticides. This, in turn, should prolong the effective life of such control agents.

### Eliminating Fly Breeding Sites:

Certain conditions around cattle, hog, poultry, and horse operations encourage flies and should be brought under control or eliminated as an aid to fly control. These include:

- For chickens, removing broken eggs and dead birds.
- Cleaning up of feed spills, manure spills, especially if wet.
- Reducing feed spills in the manure pits.
- Reducing moisture in manure in the pits.
- Repairing water leaks that cause wet manure.
- Cleaning out weed-choked water drainage ditches.
- Minimizing sources from other fly-infested animal operations in close proximity to the poultry house or livestock pens.

### How to Use NEPOREX 2SG

NEPOREX 2SG is a larvicide with specific and unique properties developed for the control of fly larvae in manure and other breeding sites. It can be used simultaneously or alternatively with fly adulticides and is thus an important component of any integrated fly control program.

NEPOREX 2SG acts as a growth regulator and affects the molting process of fly larvae. It does not kill adult flies. The elimination of adult fly population occurs gradually and becomes visible one to two weeks after application. To achieve a faster reduction of the adult fly population, the concurrent use of an adulticide is recommended.

NEPOREX 2SG is a water-soluble granule suitable for dry scattering or spraying. The application rate is 0.02 lb cyromazine per 200 sq. ft., independent of the method of application. Apply NEPOREX 2SG only on fly breeding sites. It is not intended for application to fly resting sites (walls, ceilings, etc.), as it has no effect on adult flies. Prevent contamination of feed troughs and water troughs.

Application Method	Neporex 2SG Rate Per 200 Sq. Ft.
Dry scattering	1 lb
Spraying	1 lb per gallon of water

**Dry scattering:** Apply NEPOREX 2SG directly and evenly on the fly breeding sites. Use dry scattering only in case of wet or liquid manure. Wear gloves when scattering

**Spraying:** Thoroughly mix the indicated quantity of NEPOREX 2SG with the corresponding volume of water. Apply as a course low-pressure spray on fly breeding sites with hand-pressurized or power-operated sprayers, or other suitable spray equipment.

### Note

Do not administer via feed or drinking water or contaminate feed troughs or water troughs.

### Treatment Interval

The recommended treatment interval depends on management and housing systems as well as on climatic conditions. It can vary from 21 days up to several months. More specific recommendations are given in the application schedules below. Do not apply NEPOREX 2SG more frequently than once every 21 days.

### Notes

- (1) DO NOT APPLY NEPOREX 2SG DIRECTLY TO LIVESTOCK OR LIVESTOCK FEED AS ILLEGAL RESIDUES MAY RESULT.
- (2) DO NOT FEED MANURE TREATED WITH NEPOREX 2SG TO ANIMALS.
- (3) TO AVOID ILLEGAL RESIDUES, ALLOW 1 DAY (24 HOURS) BETWEEN LAST APPLICATION AND SLAUGHTER.
- (4) DO NOT USE NEPOREX 2SG IN CONJUNCTION WITH LARVADEX 1% PREMIX IN CHICKENS. IF CHICKENS HAVE BEEN FED LARVADEX-TREATED FEED, DO NOT APPLY NEPOREX 2SG TO MANURE.
- (5) MANURE TREATED WITH NEPOREX 2SG MAY BE USED AS A SOIL FERTILIZER SUPPLEMENT. DO NOT APPLY MORE THAN 4 TONS OF MANURE TREATED WITH NEPOREX 2SG PER ACRE PER YEAR. DO NOT APPLY TREATED MANURE TO SMALL GRAIN CROPS THAT WILL BE HARVESTED OR GRAZED, OR ILLEGAL RESIDUES MAY RESULT.

### Application Schedules

#### Cattle Facilities

**Deep litter:** Treat edges and spillage areas around feed troughs and water troughs and where manure accumulates. In calf hutches placed outside, treat the whole floor surface. Apply NEPOREX 2SG by spraying approximately 3 days after manure removal and repeat treatment after each removal or when fly larvae are found.

**Slatted floor:** Treat areas where manure and other refuse accumulates to create a fly breeding site where fly larvae are present.

#### Hog Facilities

**All-in, all-out system:** Apply NEPOREX 2SG preferably by scattering immediately after cleaning and prior to the installation of the pigs on any fly breeding sites where fly larvae are present.

**Slatted floors:** Treat areas where manure or other refuse accumulates to create a fly breeding site where fly larvae are present.

**Deep litter:** Apply NEPOREX 2SG by spraying approximately 3 days after manure removal and repeat treatment after each removal or when fly larvae are found. Treat entire manure surface and wet spots around feed troughs and water troughs.

#### Poultry Facilities

Apply NEPOREX 2SG by spraying the entire manure area approximately 1 week after manure removal. Repeat treatment after each removal or when fly larvae are found.

#### Horse Barns (for horses not intended for slaughter or food)

Stalls, manure pile: Treat all areas where manure, spilled feed, and/or moisture accumulate. Scatter or spray treatment sites. Scatter NEPOREX 2SG underneath bedding on stall floors after cleaning. Repeat treatment after each removal or when fly larvae are found.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Storage:** Store in a cool, dry place. Do not store this product under wet conditions. Avoid direct sunlight and temperatures exceeding 95° F (35° C).

601259 USA – s 922739

Neporex 2 SG – 11 pounds (5 kg) – 350x125 mm – 601259 USA – Cyan  
Neporex 2 SG – 11 pounds (5 kg) – 350x125 mm – 601259 USA – Magenta  
Neporex 2 SG – 11 pounds (5 kg) – 350x125 mm – 601259 USA – Gelb  
Neporex 2 SG – 11 pounds (5 kg) – 350x125 mm – 601259 USA – Schwarz