

Oxytetracycline Injection 200 (oxytetracycline) is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline. Each mL contains 200 mg of oxytetracycline base as oxytetracycline amphoteric, and on a w/v basis, 40.0% 2-pyrrolidone, 5.0% povidone, 1.8% magnesium oxide, 0.2% sodium formaldehyde sulfoxylate (as a preservative), monoethanolamine and/or hydrochloric acid as required to adjust pH.

CAUTION:

When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

WARNINGS:

Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food.

PRECAUTIONS:

Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly or subcutaneously per injection site in adult beef and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

010670L0?

TAKE TIME



OBSERVE LABEL DIRECTIONS

Lot No.:

Exp. Date:

NDC 55529-004-05

Oxytetracycline Injection 200

(oxytetracycline)

200 mg/mL
ANTIBIOTIC

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline. **For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.**

ANADA 200-306, Approved by FDA

Net Content: 500 mL



Norbrook®



DOSAGE:

CATTLE: A single dosage of 9 mg of oxytetracycline per lb of body weight (4.5 mL/100 lb) administered intramuscularly or subcutaneously is recommended in the treatment of the following conditions:

- (1) bacterial pneumonia caused by *Pasteurella* spp. (shipping fever) in calves and yearlings, where retreatment is impractical due to husbandry conditions, such as cattle on range, or where repeated restraint is inadvisable.
- (2) infectious bovine keratoconjunctivitis (pink eye) caused by *Moraxella bovis*.

SWINE: A single dose of 9 mg of oxytetracycline per lb of body weight (4.5 mL/100 lb) administered *intramuscularly* is recommended in the treatment of bacterial pneumonia caused by *Pasteurella multocida* in swine, where retreatment is impractical due to husbandry conditions or where repeated restraint is inadvisable.

Refer to Package Insert for Complete Directions.

STORAGE:

Store at 59° to 86°F (15° to 30°C). Keep from freezing.

Not for Human Use. For Animal Use Only.

Restricted Drug - California. Use Only as Directed.

Made in the UK.

Norbrook Laboratories Limited,
Newry, BT35 6PU, Co. Down, Northern Ireland.