

# Pennchlor<sup>®</sup> 100 G<sup>™</sup>

## Chlortetracycline Type A Medicated Article

<b>Active Drug Ingredients:</b>		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 100 grams Chlortetracycline Hydrochloride/lb.	
<b>Ingredients:</b>		Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.	
<b>For Use in the Manufacture of Medicated Feeds</b> <b>CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS</b>			
<b>Directions For Use</b>			
<b>Indications For Use</b>	Use Levels of Chlortetracycline	lbs. of Pennchlor 100 G <sup>®</sup> per ton	<b>Indications For Use</b>
<b>Chickens</b>			<b>Swine</b>
<p><b>For Chickens:</b> Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)  <b>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Chickens:</b> Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)  <b>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Chickens:</b> Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5 days)  <b>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.</b></p>	100-200g/ton	1.0-2.0	<p><b>For Swine:</b> Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by <i>Group E Streptococci</i> susceptible to chlortetracycline.</p> <p><b>For Breeding Swine:</b> Control of leptospirosis (reducing the instances of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)</p> <p><b>For Swine:</b> Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Feed for not more than 14 days)  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p>
<b>Turkeys</b>			<b>CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE</b> <b>WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR THIS PRODUCT IN PRE-RUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.</b>
<p><b>For Turkeys:</b> Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)  <b>WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Turkeys:</b> Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days)  <b>WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Turkeys:</b> Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.</p> <p><b>For Turkeys:</b> Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days)  <b>WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</b></p>	200g/ton	2.0	<p><b>For Growing Cattle (over 400 lbs.):</b> For the reduction of the incidence of liver abscesses.  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Beef Cattle:</b> For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline.  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Beef Cattle (under 700 lbs.):</b> Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Beef Cattle (over 700 lbs.):</b> Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p> <p><b>For Calves, Beef, and Nonlactating Dairy Cattle:</b> For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Treat for not more than 5 days)  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p>
<b>Sheep</b>			
<p><b>For Breeding Sheep:</b> Reducing the incidence of (vibronic) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.  <b>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</b></p>	80 mg/head/day		
Approved by FDA under NADA # 138-935 Pennchlor 100 G <sup>®</sup> is a registered trademark of Pharmgate Inc.			