PharmGate Animal Health

Pennox 200 *Hi-Flo*®

Pennox 200 *Hi-Flo*®

Oxytetracycline
Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

See mixing directions, claims, cautions and warnings on back.
Restricted Drug (California) - Use Only as Directed.
Not For Human Use.

Net wt. 50 LB (22.68 Kg)

PharmGate Animal Health

PharmGate Animal Health

Pennox 200 *Hi-Flo*®

Pennox 200 *Hi-Flo*®

Oxytetracycline
Type A Medicated Article

Active Drug Ingredients	Oxytetracycline (from oxytetracycline Hydrate base) equivalent to 200 grams Oxytetracycline Hydrochloride/lb.
Ingredients	Oxytetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.

For Use in the Manufacture of Medicated Feeds. Do Not Feed Undiluted.
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS

Directions For Use

Indications For Use	Use Level of Oxytetracycline	Use of PENNOX 200 HiFlo per ton
Calves		
For Chickens: Control of infections susceptible to oxytetracycline: Control of bacterial enteritis caused by Mycoplasma synoviae: control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline. (Feed continuously for 7-14 days) RESIDUE WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO DAY WITHDRAWAL PERIOD. DO NOT USE IN LOW CALCIUM FEED CONDITIONS (LESS THAN 0.85% DIETARY CALCIUM. USE INCREASED MAY RESULT IN VIOLATIVE RESIDUES.	100-200g/ton	02-10
For Turkeys: Control of chronic respiratory disease (CRD) caused or aggravated by Mycoplasma gallisepticum and Pasteurella multocida susceptible to oxytetracycline. (Feed continuously for 7-14 days) RESIDUE WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO DAY WITHDRAWAL PERIOD. DO NOT USE IN LOW CALCIUM FEED CONDITIONS (LESS THAN 0.85% DIETARY CALCIUM. USE INCREASED MAY RESULT IN VIOLATIVE RESIDUES.	400g/ton	20
For Broiler Chickens: Reduction of mortality due to air sacculitis or infection caused by Escherichia coli susceptible to oxytetracycline. (Feed continuously for 5 days) RESIDUE WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO DAY WITHDRAWAL PERIOD. DO NOT USE IN LOW CALCIUM FEED CONDITIONS (LESS THAN 0.85% DIETARY CALCIUM. USE INCREASED MAY RESULT IN VIOLATIVE RESIDUES.	500g/ton	25
Turkeys		
For Turkeys: Control of bacterial enteritis caused by Clostridium perfringens susceptible to oxytetracycline. (Feed continuously for 5-7 days) RESIDUE WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.	300g/ton	05
For Turkeys: Control of coryza (infectious sinusitis) caused by Mycoplasma gallisepticum susceptible to oxytetracycline. (Feed continuously for 7-14 days) RESIDUE WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.	200g/ton	10
For Turkeys: Control of coryza (infectious sinusitis) caused by Haemophilus paragallinarum susceptible to oxytetracycline. (Feed continuously for 7-14 days) RESIDUE WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.	25 mg/lb body weight/day	
Sheep		
For Sheep: Treatment of bacterial enteritis caused by Escherichia coli and Clostridium perfringens and Pasteurella multocida susceptible to oxytetracycline. (Feed continuously for 7-14 days) RESIDUE WARNING: DO NOT FEED TO FEMALE SHEEP 60 DAYS OR MORE OF AGE.	10mg/lb body weight/day	
Approved by FDA		

Pennox 200 Hi-Flo® is the registered trademark of Pharmgate LLC.

Indications For Use	Use Level of Oxytetracycline	Use of PENNOX 200 HiFlo per ton
Swine		
For Swine: Treatment of bacterial enteritis caused by Erlichia coli and Salmonella choleraesuis susceptible to oxytetracycline. (Feed continuously for 5-7 days) prior to the outset of Pasteurella multocida susceptible to oxytetracycline. (Feed continuously for 7-14 days)	8mg/kg approximately 400g/lb. varying with body weight and feed consumption to provide 10mg/lb body weight/day	
For Breeding Swine: Control and treatment of leptospirosis including the incidence of abortion and shedding of leptospirae (caused by Leptospira pomona serogroup) susceptible to oxytetracycline. (Feed continuously for 30 days) RESIDUE WARNING: ZERO DAY WITHDRAWAL PERIOD.	8mg/kg approximately 400g/lb. varying with body weight and feed consumption to provide 10mg/lb body weight/day	
Calves, Beef Cattle, and Nonlactating Dairy Cattle		
RESIDUE WARNING: THIS DRUG PRODUCT IS NOT APPROVED FOR USE IN FEMALE DAIRY CATTLE 20 MONTHS OF AGE OR OLDER, INCLUDING DRY DAIRY COWS. DO NOT USE IN CATTLE TO BE SLAUGHTERED OR GIVEN WITHIN 48 HOURS, AND DO IN CALVES BORN TO THESE COWS.		
For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses. (Use continuously)	75 mg/head/day	
For Cattle: For the prevention and treatment of the incidence of Anaplasmosis. (Feed continuously) (Feed 0-5 days before and after anticipated exposure)	02-20g/head/day	
For Calves, Beef Cattle and Nonlactating Dairy Cattle: Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and liver complicated caused by Pasteurella multocida susceptible to oxytetracycline. (Feed continuously for 5-14 days) RESIDUE WARNING: 5 DAYS WITHDRAWAL PERIOD AT SINGLE DOSAGE.	10 mg/lb body weight/day	
Honey Bees		
For Honey Bees: For control of American Foulbrood caused by Paenibacillus larvae and European Foulbrood caused by Streptococcus pluton susceptible to oxytetracycline. RESIDUE WARNING: REMOVE AT LEAST 4 WEEKS PRIOR TO MAIN NECTAR FLOW. SUBSTITUTION AVAILABILITY ON REQUEST	200mg/colony	

PharmGate Animal Health

Marketed by:
Pharmgate Animal Health, LLC
1040 Industrial Road
Omaha, Nebraska 68114

19-29 10-02
Rev 01/15
Pag. 1