

IN ANIMAL REARING AND CONFINEMENT FACILITIES

TO DISINFECT COMMERCIAL ANIMAL CONFINEMENT FACILITIES SUCH AS POULTRY HOUSES, SWINE PENS, CALF BARN AND KENNELS

It is a violation of federal law to use this product in a manner inconsistent with its labeling

- 1) Remove all animals and feed from premises, vehicles, enclosures, coops and crates.
- 2) Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
- 3) Empty all troughs, racks and other feeding and watering applications.
- 4) Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5) Preparation of active disinfecting solution: (500 ppm ClO₂) Place 1 1/3 fl. oz. of ProOxine® concentrate per gallon of working solution into a clean, plastic pail and add ten (10) grams (2 teaspoons) of Bio-Cide Activator Crystals or food grade citric acid of no less than 99% purity. Prepare in a well ventilated area; avoid breathing any fumes which may be produced while crystals are dissolving. Allow five (5) minutes reaction time for crystals to dissolve completely. To this solution, add one (1) gallon of clean potable water. This will yield a working solution containing 500 ppm of available chlorine dioxide.
- 6) To apply: Using a commercial sprayer, saturate all surfaces with the activated ProOxine® solution for a period of ten (10) minutes. Active solution may be irritating when breathed, therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate only when solution has dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.

TO DISINFECT POULTRY HOUSES

Special Instructions for Inactivating Avian Influenza A Kills Avian Influenza A on pre-cleaned environmental surfaces

- 1) Remove all poultry and feed from premises, vehicles, enclosures, coops and crates.
- 2) Remove all litter and droppings from floors, walls and surfaces of facilities and fixtures occupied or traversed by poultry.
- 3) Empty all troughs, racks and other feeding and watering applications.
- 4) Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5) Preparation of active disinfecting solution: (500 ppm ClO₂) Place 1 1/3 fl. oz. of ProOxine® concentrate per gallon of working solution into a clean, plastic pail and add ten (10) grams (2 teaspoons) of Bio-Cide Activator Crystals or food grade citric acid of no less than 99% purity. Prepare in a well ventilated area; avoid breathing any fumes which may be produced while crystals are dissolving. Allow five (5) minutes reaction time for crystals to dissolve completely. To this solution, add one (1) gallon of clean potable water. This will yield a working solution containing 500 ppm of available chlorine dioxide.
- 6) To apply: Using a commercial sprayer, saturate all surfaces with the activated ProOxine® solution for a period of ten (10) minutes. Active solutions may be irritating when breathed, therefore, always use an applicable NIOSH/MSHA approved respirator appropriate for chlorine dioxide when spraying these solutions. Immerse all equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and droppings. After treatment, ventilate buildings, coops or other enclosed spaces and allow to air dry. Repopulate only when solution has dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before use.

TO CONTROL THE BUILD-UP OF ODOR AND SLIME FORMING BACTERIA IN ANIMAL CONFINEMENT AREAS

- 1) Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, cases and other facilities and fixtures occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with clean water.
- 2) Preparation of solution: Place 2.6 fl. oz. ProOxine® concentrate per gallon of working solution (1000 ppm available ClO₂) into a clean, plastic pail. Dilute concentrate with one (1) gallon clean, potable water.
- 3) To apply: Using a commercial sprayer or fogger, saturate all surfaces with the ProOxine® solution.

TO DISINFECT WATER SUPPLY FOR POULTRY, SWINE, CATTLE AND OTHER LIVESTOCK

- i) Use Bio-Cide International, Inc. automated activation equipment to generate an aqueous chlorine dioxide solution. Alternatively, ProOxine® may be manually activated to generate an aqueous chlorine dioxide solution. The activated ProOxine® solution can be either batch loaded or metered into the poultry and drinking water supply at a point in the system which insures uniform mixing and distribution of up to 5 ppm chlorine dioxide.
- ii) Automated Activation Equipment Method: Bio-Cide International, Inc. automated activation equipment may be used to generate an aqueous chlorine dioxide solution for metering into the water supply to treat 3 to 5 ppm activated ProOxine® concentration.
- iii) Manual Activation Method: Activated ProOxine® concentrate may be prepared by manual mixing and subsequent dilution for treatment of the water supply at 3 to 5 ppm according to the activation and dilution charts. For example, to manually prepare activated ProOxine® to treat 1,000 gallons of water at 3 ppm activated ProOxine® :

1) Preparation of active solution: Place 8 fl. oz of ProOxine® concentrate into a plastic container and add 60 grams of citric acid. Prepare in a well ventilated area. Avoid breathing and fumes while crystals are dissolving. Gently stir until citric acid crystals are completely dissolved. Allow five minutes reaction time.

2) The activated concentrate may then be added to 1,000 gallons of water. Allow ten minutes before delivery to livestock water lines.

TO DISINFECT POTABLE WATER

For most municipal and other potable water systems, a chlorine dioxide residual concentration up to 2.0 ppm is sufficient to provide adequate disinfection. Typically, the target residual concentrations range from 0.20 - 0.75 ppm. Monitor the distribution system to ensure that the chlorite concentration does not exceed its maximum contaminant level (MCL) of 1 mg/L and that chlorine dioxide does not exceed its maximum residual disinfection level (MRDL) of 0.8 mg/L. For wastewater and sewage applications, residual chlorine dioxide concentrations up to 5.0 ppm are generally adequate.



ProOxine®

SANITIZER DISINFECTANT Bacteriostat/Deodorizer

FUNGICIDAL - BACTERICIDAL - VIRUCIDAL
ANIMAL REARING AND CONFINEMENT FACILITIES

ProOxine has demonstrated efficacy
against the following animal viruses:

Avian Influenza A Virus

Porcine Epidemic Diarrhea Virus

Porcine Reproductive & Respiratory Disease Syndrome Virus

Pseudorabies Virus

Newcastle Disease Virus

Canine Parvovirus

Low Corrosion - Guaranteed Shelf Life - Non-Flammable

Active Ingredient:
Chlorine Dioxide.....5.0%
Other Ingredients.....95.0%
Total100.0%

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

CAUTION

STORE IN A DARK PLACE

E.P.A. REG # 9804-9

KEEP FROM FREEZING

E.P.A. Est. No. 9804-OK-1



Bio-Cide International Inc.
Norman, OK 73070
www.bio-cide.com

Net Contents: 5 gallon

TO CONTROL THE BUILDUP OF SLIME AND ODOR CAUSING BACTERIA IN FEED WATERS

- 1) Feed water should be treated at the rate of 0.4 fl. oz. of ProOxine® per 30 gallons of water (5ppm available ClO₂) and may be injected or batch loaded.
- 2) Feed water storage tanks should be sufficiently sealed to prevent outside contamination and direct sunlight.

TO SANITIZE CLEAN SHELL EGGS INTENDED FOR FOOD OR FOOD PRODUCTS

- 1) Preparation of sanitizing solution: Prepare an activated working solution containing 100-200 ppm available chlorine dioxide.
- 2) Spray eggs thoroughly with activated solution making sure surface area is thoroughly wet for at least one (1) minute and allow to drain. Solution must be equal to or warmer than the eggs, but not to exceed 130° F.
- 3) Eggs that have been sanitized with this chlorine dioxide compound may be broken in the manufacture of egg products without a prior potable water rinse. Eggs must be reasonably dry before casing or breaking. Never reuse activated solutions.

STORAGE AND DISPOSAL

Product Storage: Store in a cool dark area in original container. Avoid storage in direct sunlight. In case of spill, flood with water before discarding to drain. Do not contaminate water, food, or feed by storage or disposal.

Container Disposal: Do not reuse or refill this container. Offer for recycling if available. Offer for reconditioning if appropriate. Triple rinse container, or equivalent, promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for ten seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this process two more times.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals: CAUTION: Causes moderate eye irritation. Harmful if swallowed. Harmful if inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or going to the restroom. Remove contaminated clothing and wash before reuse. Handlers applying chlorine dioxide in an occupational setting must wear gloves.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates, oysters and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or public waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA

FIRST AID

If in eyes	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lens, if present, after the first 5 minutes then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
If swallowed	- Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by a poison control center or doctor. - Do not give anything by mouth to an unconscious person.
If inhaled	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. - Call a poison control center or doctor for treatment advice.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24 hour emergency information on this product call NPIC at 800.858.7378

For 24 hour transportation emergency information on this product call Chemtrec at 1-800-424-9300 (U.S., Canada, Puerto Rico, Virgin Islands) or 703-527-3887 (All Other Areas)

Seller expressly warrants that the product conforms to its chemical description. To that extent consistent with applicable law, seller makes no other warranties, either express or implied, including, but not limited to, the warranties of merchantability or fitness for a particular purpose or use, that extend beyond the statements made on this label.

(rev 12-22-15)